

## A NOTE ON THE KERNEL ESTIMATION OF ATKINSON'S INDEX OF INEQUALITY

K. AGBOKOU\*, K.M. MIDAGBODJI

Received: 03.12.2024 / Revised: 20.02.2025 / Accepted: 27.02.2025

**Abstract.** *World Bank macrodata of every country on our planet show that national incomes per capita account for a major part of population disparity and these follow usual distributions well known in the literature in almost all continents. Measuring and comparing disparity is a huge task that requires assembling the relative nature of small and large national incomes without distinctions. This is the main reason that pushes us to consider in this paper the Atkinson inequality index (continuous case) developed towards the end of the 20th century in order to measure this disparity. Since then its nonparametric estimator has not been developed but rather a well-known classical discrete form. This makes the estimation or measurement of economic inequalities not very complex. In this paper, we construct a kernel estimator of the Atkinson inequality index and by extension that of its associated welfare function, then we establish their almost sure asymptotic convergences. Finally, we explore the performance of our estimators in a simulation study and draw conclusions about national incomes per capita on each continent and then globally by making comparisons with the classical form, based on World Bank staff estimates based on sources and methods from "The Changing Wealth of Nations". The results obtained highlight the interest of kernel-based measures as well as the sensitivity of the index with respect to the inversion parameter.*

**Keywords:** probability density, kernel estimation, statistical inference, likelihood, welfare

**Mathematics Subject Classification (2020):** 60E05, 60F05, 62F10, 62G05, 62G20

### 1. Introduction

Measuring inequality is essential to understanding the distribution of wealth in a society and changes in the social structure. It helps guide public policies, particularly those of

---

\* Corresponding author.

**Komi Agbokou**

University of Kara, Kara, Togo  
E-mail: ffomesteino@yahoo.com

**Koami Mawuko Midagbodji**

University of Lomé, Lomé, Togo  
E-mail: midakoma@gmail.com

redistribution, and measure their impact. To measure economic inequality, there is a wide variety of tools that represent different points of view on the object studied. After reflecting on the relevant level of analysis of inequality, this article presents the different indicators that allow us to assess the extent of economic inequality, its evolution and its persistence. It then shows the normative scope of the choice of indicators in relation to considerations of social justice. After having been ignored for several decades, the issue of inequality is back in the spotlight. The recent reopening of inequality raises questions and arouses serious concerns. Inequalities refer to differences that generate phenomena of social hierarchy. Differences can relate to the allocation of a resource that is unequally distributed or refer to unequal access to certain goods or services: this is how we speak of income inequality. Economic inequalities traditionally refer to inequalities of income and wealth. These inequalities are the subject of particular attention due to the importance in our societies of the economic dimension in the social valuation of individuals. If not all inequalities are unfair, the assessment of the fair or unfair nature of a situation is always done in the light of a standard of equality in relation to which the situation is evaluated. Thus, depending for example on whether we value equality of situations or equality of rights, a different view will be taken of the same situation. It is thus impossible to think of inequalities without reference to a conception of social justice in relation to which a judgment will be made on reality. This renewed interest in the study of inequalities stems in part from publications that have highlighted their increase. We can notably cite the work of Piketty [10] on this question. One of the main contributions of this author (with others) is to have enabled the dissemination of statistics, in particular on high incomes, making it possible to precisely measure the evolution of inequalities. Documenting the existence of inequalities and having precise information on their variation is a necessary prerequisite for any scientific debate. Far from being simply technical, the discussion around the choice of a relevant indicator to measure inequalities is in reality a scientific challenge.

It should also be noted that the issue of the well-being of the population is of major importance in the political, social and economic spheres. Well-being means that the population has sufficient means to meet its needs, organise its life independently, use and develop its abilities and pursue its goals. This cannot happen without appropriate framework conditions. The term well-being is used here as a synonym for quality of life. Well-being is considered not only in its material and financial dimensions, but in a broader perspective that encompasses the immaterial situation of the population. Material resources are income and wealth, which enable the individual to meet his or her needs. But other material dimensions, such as housing and work, are taken into account in measuring well-being. Education, health and social relations are part of the immaterial dimensions of well-being, which also encompass the legal and institutional framework that allows citizens to participate in political life and ensures the physical security of people. Finally, the concept of well-being includes environmental aspects such as water quality, air quality and noise pollution. In an approach to well-being that aims to be as broad as possible, it is important to consider not only objective living conditions, but also their subjective perception by the population, namely what the population thinks, for example, of its housing conditions and the state of the environment, its feeling of security, its degree of satisfaction with the world of work and with life in general. In-

equality indices aim to provide information on the situation of the population. To this end, it is necessary to grasp a large number of elements that constitute well-being and to describe its different facets. Inequality indices provide statistical information on the state and evolution of well-being in a broad context, and which can serve as a basis for forming public opinion and making political decisions (see also Harper [14]).

Inequality indices are classified by categories. In this paper, we will cite the most widespread ones. Generally speaking, inequality measures fall into two broad categories depending on the approach used to calculate them: descriptive measures and normative measures (see Sen [18]). Descriptive inequality measures are usually mathematical or statistical formulas, for example the Gini index (see Agbokou et al. [5] and also Banerjee [8]). Therefore, the characteristics of these indices depend on their mathematical or statistical properties, respectively. Most inequality indices are descriptive in nature. Normative inequality indices are derived from a social welfare function based on a prior value judgment about the effects of inequality on social welfare. These measures combine the inequality index with a social evaluation and specify whether inequality is harmful or not, as well as the degree of welfare that a society loses or gains because of this inequality. Atkinson [6], [7] inequality indices are among the most frequently cited normative measures. Note that the inequality indices examined here do not necessarily satisfy all of axioms (see Sen et al. [19]). For example, the Atkinson index satisfies almost all of the axioms, but it is not additively decomposable. Note also that there are also inequality measures derived from entropy that we will not discuss here.

We denote by  $X$  a continuous random variable representing income. For simplicity, let  $X$  be strictly positive (no zero or negative income),  $f(x)$  is its probability density,  $F(x)$  is its distribution function representing the cumulative distribution of income and  $\mu$  the average income. The consideration of a welfare function to define an inequality index is illustrated first and foremost by the Atkinson index. We start with the following welfare function:

$$W(\theta) = \int_{\mathbb{R}} \frac{x^{1-\theta}}{1-\theta} f(x) dx. \quad (1)$$

The inequality measure that we deduce from this simple welfare function is written as:

$$A(\theta) = 1 - \left[ \int_{\mathbb{R}} \left( \frac{x}{\mu} \right)^{1-\theta} f(x) dx \right]^{\frac{1}{1-\theta}}, \quad (2)$$

where  $\theta \geq 0$  is the inequality aversion parameter. We generally use values between 0 and 1 for  $\theta$ . For  $\theta = 1$ , the form is indeterminate and we remove the indeterminacy by taking as a welfare function:

$$W(1) = \int_{\mathbb{R}} \log(x) f(x) dx,$$

and the corresponding form of the index in this case is therefore

$$A(1) = 1 - \frac{1}{\mu} \exp \left[ \int_{\mathbb{R}} \log(x) f(x) dx \right]. \quad (3)$$

For  $\theta \rightarrow 0$ , we come across a Rawlsian measure, Rawls [17], where only the fate of the poorest matters to society. The Atkinson index has a value in  $[0, 1]$ . It is worth 1 when an


individual has everything and the others nothing. The welfare function associated with this index is the one that weights the observations by their rank. It is the poorest who will receive the greatest weight. We therefore deduce it, according to the formulas (1) and (2), in the form:

$$W(\theta) = \begin{cases} [\mu(1 - A(\theta))]^{1-\theta}, & \text{if } \theta \neq 1, \\ \log[\mu(1 - A(\theta))], & \text{if } \theta = 1. \end{cases} \quad (4)$$

Several authors have worked on the parametric estimation of the Atkinson index. We can start by citing Guerrero [13] who was one of the first to estimate the aversion parameter to determine the degree of inequality for a given income distribution, with a construction of confidence intervals. We can also cite Biewen et al. [10] and Tchamyou [21] to name only these. The aim of this paper is to provide a nonparametric estimation of the Atkinson inequality index (2), based on the kernel method. Then a simulation study will be done on simulated or real data. Finally we will make a comparative study of this estimator with the one that already exists in the literature.

## 2. Description, Materials and Methods

### 2. 1. Interpretation of the Atkinson index: graphical approach and approximation



**Fig. 1. Atkinson inequality index graphic description**

The basis of the measurement of Atkinson inequalities is based on the concept of equitably distributed equivalent income that we denote  $X^*$ . Income  $X^*$  is the income that, if all individuals had this amount, would give the same level of social utility as the existing one ( $\bar{W}$ ). Taking the case of two individuals, we can graphically represent the construction of the Atkinson index. Figure 1 illustrates this concept of equitably distributed equivalent. This graph shows the welfare function constructed on the space of individual incomes. The ( $Ox$ ) axis shows the income of individual 1, while the ( $Oy$ ) axis shows that of individual 2. If  $X_0$  is the income of the first individual and  $X_1$  that of the second, the average income is  $\bar{X}$ . Suppose the income distribution is such that point A prevails i.e. the point where we have  $X_0 > X_1$ . In the absence of inequality aversion ( $\theta = 0$ ), utilitarian welfare would prevail i.e. the straight line. With this welfare function, the only way to have equal incomes at the same level of welfare is therefore to give an average income to both individuals i.e.  $\bar{X}$ . Since inequality aversion is zero, we

are not willing to reduce the size of the butter to have more equal shares. With inequality aversion, the convex welfare function would prevail. Now, starting from  $A$ , we can find a point where incomes are equally distributed at the same level of welfare. Since the welfare function is convex, income  $X^*$  must be less than average income  $\bar{X}$ . Income  $X^*$  is the abscissa of point  $B$  of the 45 degree line that has the same social welfare as  $A$  and  $C$ . Even though total income (the sum of the two individual incomes) is less than  $X_A$ , it is compensated by the gain in equality of the distribution. The reason being that, since inequality aversion is positive, we are now willing to pay the lower price of butter in order to have more equal shares. Equality is measured by the ratio  $X^*/\bar{X}$ . If this ratio is equal to 1 then each individual has the same level of income or if the welfare function is utilitarian (there is no perceived inequality). The approximatif Atkinson inequality index can therefore be expressed as follows (see Bellù et al. [9]):

$$\tilde{A} = 1 - \frac{X^*}{\bar{X}}.$$

Instinctively, this index tells us how much income we are willing to give up to have equal income. We therefore deduce that if the welfare is of the discrete form

$$\tilde{W}(\theta) = \begin{cases} \frac{1}{n} \sum_{i=1}^n \frac{X_i^{1-\theta}}{1-\theta}, & \text{if } \theta \neq 1, \\ \frac{1}{n} \sum_{i=1}^n \log(X_i), & \text{if } \theta = 1, \end{cases} \quad (5)$$

then the discrete Atkinson index is given by:

$$\tilde{A}(\theta) = \begin{cases} 1 - \left[ \frac{1}{n} \sum_{i=1}^n \left( \frac{X_i}{\bar{X}} \right)^{1-\theta} \right]^{\frac{1}{1-\theta}}, & \text{if } \theta \neq 1, \\ 1 - \frac{1}{\bar{X}} \exp \left[ \frac{1}{n} \sum_{i=1}^n \log(X_i) \right], & \text{if } \theta = 1. \end{cases} \quad (6)$$

The formulas (or estimators if we can say so) (5) and (6) are the most used in the literature so far and would reveal inadequacies following the variation of the data since most of the data follow a distribution or at worst approximately a distribution. Hence the need to construct a non-parametric estimator or kernel estimator which often presents satisfactory results.

## 2. 2. Construction of the kernel estimator

Kernel estimation (or Parzen-Rosenblatt method or KDE) is a non-parametric method for estimating the probability density of a random variable. It is based on a sample of a statistical population and makes it possible to estimate the density at any point of the support. In this sense, this method cleverly generalizes the histogram estimation method. Kernel density estimation is used to obtain a smooth estimate of the data distribution

without making assumptions about the shape of the distribution and it is famous for its surprising smoothing character.

Let  $(X_i)_{1 \leq i \leq n}$  be a random sample of size  $n$  from a population  $X$  with density function  $f$ , which represents the distribution of income.  $X_i$ , for  $i \in [1, n]$ , designating the respective income of the  $n$  individuals, are independent and identically distributed (i.i.d.) observations. The main of nonparametric density estimation is to estimate  $f$  with as few assumptions about  $f$  as possible. Among the existing kernel density estimators, we choose the classic and most famous case well known in the literature, because of its simplicity for independent observations. This estimator depends on a parameter called smoothing parameter or bandwidth parameter. It is defined by:

$$\widehat{f}_n(x) = \frac{1}{nh} \sum_{i=1}^n K\left(\frac{x - X_i}{h}\right), \quad x \in \mathbb{R}, \quad (7)$$

where  $h = h(n) = h_n$  is the bandwidth parameter depending on the sample size  $n$  and  $K$  a probability density function called kernel function verifying certain particular properties called regularity properties.

From the estimator (7) of the distribution function  $f$ , it is obvious that we obtain an estimator of the Atkinson index (2), which is defined by:

$$\begin{aligned} \widehat{A}_n(\theta) &= 1 - \left[ \int_{\mathbb{R}} \left(\frac{x}{\widehat{\mu}_n}\right)^{1-\theta} \widehat{f}_n(x) dx \right]^{\frac{1}{1-\theta}} = \\ &= 1 - \left[ \frac{1}{nh} \sum_{i=1}^n \int_{\mathbb{R}} \left(\frac{x}{\widehat{\mu}_n}\right)^{1-\theta} K\left(\frac{x - X_i}{h}\right) dx \right]^{\frac{1}{1-\theta}}, \end{aligned} \quad (8)$$

if  $\theta \neq 1$ ,

and

$$\begin{aligned} \widehat{A}_n(1) &= 1 - \frac{1}{\widehat{\mu}_n} \left[ \int_{\mathbb{R}} \log(x) \widehat{f}_n(x) dx \right] = \\ &= 1 - \frac{1}{\widehat{\mu}_n} \exp \left[ \frac{1}{nh} \sum_{i=1}^n \int_{\mathbb{R}} \log(x) K\left(\frac{x - X_i}{h}\right) dx \right], \end{aligned} \quad (9)$$

where  $\widehat{\mu}_n$  is the estimator average income  $\mu$ . It is easy to verify that  $\widehat{\mu}_n$  satisfies:

$$\widehat{\mu}_n = \int_{\mathbb{R}} x \widehat{f}_n(x) dx = \frac{1}{n} \sum_{i=1}^n X_i.$$

And consequently, we can obtain the nonparametric estimator of (4)

$$\widehat{W}_n(\theta) = \begin{cases} [\widehat{\mu}_n(1 - \widehat{A}_n(\theta))]^{1-\theta}, & \text{if } \theta \neq 1, \\ \log[\widehat{\mu}_n(1 - \widehat{A}_n(\theta))], & \text{if } \theta = 1. \end{cases} \quad (10)$$

The following paragraph provides a study of the near convergence or strong consistency of our estimators and also those of some estimators which will result from them under certain regularity hypotheses.

### 3. Study of the Convergence of the Atkinson Inequality Index

In all that follows, besides the classical regularity conditions (see in Agbokou [1]), well known in the literature, on the kernel function  $K$  and on the smoothing parameter  $h$ , we consider the following hypotheses.

#### 3. 1. Assumptions

##### 3. 1. 1. The model assumptions

- $A_1$ . The random variable  $X$  takes values in  $\mathbb{R}_+$  or a compact subset of  $\mathbb{R}_+$ .  
 $A_2$ . The density function  $f$  of  $X$  is uniformly continuous on its support and satisfies

$$\int_{\mathbb{R}} x^\tau f(x) = \mathbb{E}(X^\tau) > 0, \forall \tau > 0.$$

- $A_4$ . There exists a strictly positive constant  $\eta$  such that  $\mu > \eta$  and  $\theta$  takes values in  $[0; 1[$ .  
 $A_5$ . The function  $\phi : y \mapsto y^\alpha$  on  $\mathbb{R}_+$  verifies the following conditions:  
*i.* there exist  $\kappa > 1$  strictly positive constant such:  $|\phi(u) - \phi(v)| \leq \alpha^\kappa |u - v|, \forall \alpha > 1$  and  $\forall u, v \in \mathbb{R}_+$ ;  
*ii.* there exist  $C_\alpha \geq 1$  strictly positive constant such:  $\phi(|u - v|) \leq C_\alpha |u - v|, \forall \alpha \in [0; 1]$  and  $\forall u, v \in [0; 1]$ ;  
*iii.* there exist  $Q_\alpha \geq 1$  strictly positive constant such:  $\phi(u + v) \leq Q_\alpha (u^\alpha + v^\alpha), \forall \alpha \in [0; 1]$  and  $\forall u, v \in \mathbb{R}_+$ .

##### 3. 1. 2. The kernel assumptions

$K$  is a symmetric kernel of bounded variation on  $\mathbb{R}$  vanishing outside the interval  $[-a, a]$  for some  $a > 0$  and satisfying:

- $K_1$ .  $\int_{\mathbb{R}} K(u) = 1$ ;  
 $K_2$ .  $\int_{\mathbb{R}} uK(u) = 0$ ;  
 $K_3$ .  $\int_{\mathbb{R}} u^\tau K(u) = \nu(K) > 0, \forall \tau > 0, \tau \neq 1$ .

##### 3. 1. 3. The bandwidth parameter hypothesis

The bandwidth parameter  $h = (h_n)_{n \in \mathbb{N}}$  is a sequence of positive nonincreasing real numbers satisfying:

- $H_1$ .  $h_n^\tau \rightarrow 0, \forall \tau > 0, n \rightarrow +\infty$ ;  
 $H_2$ .  $nh_n \rightarrow +\infty, n \rightarrow +\infty$ .

**Remark 1.** The assumptions  $A_1 - A_2$ ,  $K_1 - K_2$  and  $H_2$  are quite standard.  $A_1 - A_4$ ,  $A_5.i$  and  $iii.$ ,  $K_1 - K_3$  and  $H_1 - H_2$  insure the strong uniform convergence of the estimators (8) and (9) to (2) and (3) respectively. Finally all the assumptions ensure the strong uniform convergence of (10) to (4).

To simplify the writings and make them easier to handle, we adopt the following notations:

- $\Delta(\theta) = \int_{\mathbb{R}} x^{1-\theta} f(x) dx$  and  $A(\theta)$  then becomes:  $A(\theta) = 1 - \left[ \frac{\Delta(\theta)}{\mu^{1-\theta}} \right]^{\frac{1}{1-\theta}}$ ,
- $\Lambda = \exp \left[ \int_{\mathbb{R}} \log(x) f(x) dx \right]$  and  $A(1)$  is written in the form:  $A(1) = 1 - \frac{\Lambda}{\mu}$ .

Consequently, it is clear that their kernel estimators are respectively  $\widehat{\Delta}_n(\theta)$  and  $\widehat{\Lambda}_n$ .

### 3. 2. Strong consistency

In this subsection, we prove the consistency of our estimator and give a rate of convergence. Our first result is the almost sure uniform convergence with an appropriate rate of the estimators  $\widehat{\Delta}_n$  and  $\widehat{\Lambda}_n$  stated in Proposition 1, which are the key for investigating the strong consistency of  $\widehat{A}_n(\theta)$  and  $\widehat{A}_n(1)$  given by Theorem 1. The last result deals with the strong consistency of the welfare function estimator  $\widehat{W}_n(\theta)$  given by Corollary.

**Proposition 1.** *Under assumptions  $A_1$ ,  $A_5 - iii.$ ,  $K_1$ ,  $K_3$  and  $H_1$  we have:*

(i).

$$|\widehat{\Delta}_n(\theta) - \Delta(\theta)| \longrightarrow 0 \text{ a.s.}, \quad n \rightarrow +\infty \quad \forall \theta \in [0, 1[.$$

(ii). *In particular for  $\theta = 1$ , we have,*

$$|\widehat{\Lambda}_n - \Lambda| \longrightarrow 0 \text{ a.s.}, \quad n \rightarrow +\infty .$$

The Proposition 1 leads to the convergence of the inequality index function which is established by the following theorem:

**Theorem 1.** *In addition to the assumptions of the Proposition 1, if hypotheses  $A_2$ ,  $A_4 - A_5 - i.$ ,  $K_2$  and  $H_2$  are satisfied, then we obtain:*

(i).

$$|\widehat{A}_n(\theta) - A(\theta)| \longrightarrow 0 \text{ a.s.}, \quad n \rightarrow +\infty \quad \forall \theta \in [0, 1[.$$

(ii). *In particular, if  $\theta = 1$ , then we get,*

$$|\widehat{A}_n(1) - A(1)| \longrightarrow 0 \text{ a.s.}, \quad n \rightarrow +\infty .$$

As a consequence of the Theorem 1, we arrive at the strong constancy of the estimator of the welfare function (10).



**Corollary 1.** *Under the same hypotheses as Theorem 1, assume that hypothesis  $A_5 - ii$  is verified, then we obtain:*

$$|\widehat{W}_n(\theta) - W(\theta)| \longrightarrow 0 \text{ a.s.}, \quad n \rightarrow +\infty \quad \forall \theta \in [0, 1].$$

**Corollary 2.** *Under the same assumptions as Theorem 1, we assume that the sequence  $(X_i)_{1 \leq i \leq n}$  of i.i.d. random variables is such that  $\mathbb{E}(X_i) = \mu$  and  $\mathbb{V}ar(X_i) = \sigma^2 \quad \forall i \in \{1, \dots, n\}$ . Then for  $n$  large enough, we get:*

$$\widehat{A}_n(\theta) - A(\theta) \longrightarrow \mathcal{N}(0, s^2) \quad \text{a.s.} \quad \forall \theta \in [0, 1],$$

where

$$s^2 = (1 + \mu^2) \left( 1 + \frac{\sigma^2}{\mu^2} \right).$$

### 3. 3. Appendix of proofs

As for the proofs of the previous proposition, theorem and corollary, we need some very important or primordial lemmas whose demonstrations will be made or will be given here as well.

Let us first recall that if  $\{X_i\}_1^n$  are i.i.d random variables with a distribution  $F$ . Consider a parametric function  $\Pi$  for which there exists an unbiased estimator. The parametric function  $\Pi$  can be expressed in the following form:

$$\Pi(F) = \mathbb{E}[\psi(X_1, \dots, X_l)] = \int_{\mathbb{R}^l} \psi(X_1, \dots, X_l) dF(x_1) \times \dots \times dF(x_l),$$

where  $\Psi$  is a function of  $l$  i.i.d random variables from  $\{X_i\}_1^n$  with  $l \leq n$ . For any function  $\Psi$ , the corresponding U-statistic for the estimation of  $\Pi$  based on a random sample of size  $n$  is obtained by averaging  $\Psi$  symmetrically over the observations

$$U_n = U(X_1, \dots, X_n) = \frac{1}{\binom{n}{l}} \sum_s \psi(X_1, \dots, X_l),$$

where  $\sum_s$  represents the summation over the  $\binom{n}{l}$  combinations of  $l$  distinct elements  $i_j, \quad j \in \{1, \dots, l\}$  from  $\{1, \dots, n\}$ . In particular, we have:

**Lemma 1.** *For  $\psi(x) = x^m$  ( $m > 0$ ), the corresponding U-statistic is*

$$U_m = U(X_1, \dots, X_n) = \frac{1}{n} \sum_1^n X_i^m$$

and for  $n$  large enough, we get

$$U_m \longrightarrow \mu_m = \mathbb{E}[X^m] = \int_{\mathbb{R}} x^m dF(x).$$

In particular for  $m = 1$ , we set  $\widehat{\mu}_n = U_1 \longrightarrow \mu = \mu_1$ .

*Proof.* The proof of this lemma is similar to those found in Serfling [20] and Lehmann [15]. Thus we make an exception to this one. ◀

**Lemma 2.** For every  $p \in ]0; 1]$ , the function defined on  $I = [0; +\infty)$  by  $\Phi : z \mapsto z^p$  satisfies the following condition:

$$\forall u, v \in I : |\Phi(u) - \Phi(v)| \leq p|u - v|^p, \quad (11)$$

and the number  $p$  is the maximum of all values that verify this property.

*Proof.* For  $u, v$  let us suppose that  $u > v$ . Aware that  $p - 1 \leq 0$ , we have

$$\begin{aligned} |\Phi(u) - \Phi(v)| &= u^p - v^p = \int_v^u pz^{p-1} dz \\ &\leq \int_v^u p(z - v)^{p-1} dz = (u - v)^p \leq |u - v|^p \quad \text{because } u > v. \end{aligned}$$

In this case, the corresponding  $p$  of the inequality (11) which satisfies the condition is equal to 1.

On the other hand, from the relation (11), setting  $u = 1$  and  $v = 0$ . It follows  $p \geq 1$ .

Now assume that  $p > 1$ . If there exists another real number  $q$  such that  $q > p$  and verifies the inequality (11), then we obtain

$$\forall u, v \in I : |\Phi(u) - \Phi(v)| \leq q|u - v|^q. \quad (12)$$

By setting  $v = 0$  in relation (12), we have

$$|\Phi(u)| = u^p \leq u^q \quad \forall u > 0.$$

Taking into account that  $p - q < 0$ , we get  $u^{p-q} \leq 1$ . This is absurd because the left-hand side diverges towards  $+\infty$  when  $u$  tends towards 0. This brings the proof of this lemma to term. ◀

The lemmas having been established, let's move on to the proofs of the main results obtained.

*Proof. (Proposition 1)*

(i) From the sequence of functions  $\hat{\Delta}_n$  in  $\theta$ , we obtain

$$\hat{\Delta}_n(\theta) = \int_{\mathbb{R}} x^{1-\theta} \hat{f}_n(x) dx = \frac{1}{nh} \sum_{i=1}^n \int_{\mathbb{R}} x^{1-\theta} K\left(\frac{x - X_i}{h}\right) dx.$$

A change of variable, the use of hypotheses and  $A_5 - ii$ ,  $K_1 - K_3$ , allows us to write successively

$$\begin{aligned}\widehat{\Delta}_n(\theta) &= \frac{1}{n} \sum_{i=1}^n \int_{\mathbb{R}} (X_i + wh)^{1-\theta} K(w) dw , \\ &\leq \frac{Q_{1-\theta}}{n} \sum_{i=1}^n \int_{\mathbb{R}} (X_i^{1-\theta} + w^{1-\theta} h^{1-\theta}) K(w) dw , \\ &\leq Q_{1-\theta} \left[ \frac{1}{n} \sum_{i=1}^n X_i^{1-\theta} + h^{1-\theta} \int_{\mathbb{R}} w^{1-\theta} K(w) dw \right] \\ &\leq Q_{1-\theta} [U_{1-\theta} + h^{1-\theta} \nu(K)] .\end{aligned}$$

Then we get

$$|\widehat{\Delta}_n(\theta) - U_{1-\theta}| \leq Mh^{1-\theta} \text{ where } M > 0; \text{ is a constant.}$$

In other words

$$|\widehat{\Delta}_n(\theta) - U_{1-\theta}| = \mathcal{O}(h^{1-\theta}) \quad a.s.$$

Furthermore, we have

$$|\widehat{\Delta}_n(\theta) - \Delta(\theta)| \leq |\widehat{\Delta}_n(\theta) - U_{1-\theta}| + |U_{1-\theta} - \Delta(\theta)|. \quad (13)$$

For  $n$  large enough and under hypothesis  $H_1$ , Lemma 1 applied to inequality (13) completes the first part (i) of the proof of this proposition.

(ii) The second part is inspired by the first, with a few exceptions since it is a special case. Since function  $z \mapsto \exp(z)$  is a convex function, Jensen's inequality allows us to write

$$\begin{aligned}\widehat{\Lambda}_n &= \exp \left[ \int_{\mathbb{R}} \log(x) \widehat{f}_n(x) dx \right] , \\ &\leq \frac{1}{n} \sum_{i=1}^n \int_{\mathbb{R}} x K \left( \frac{x - X_i}{h} \right) dx.\end{aligned}$$

The same change of variable and the use of the same hypotheses lead to

$$\begin{aligned}\widehat{\Lambda}_n &\leq \frac{1}{n} \sum_{i=1}^n X_i + h \int_{\mathbb{R}} w K(w) dw , \\ &\leq U_1 + h\nu(K) \text{ because here } w > 0\end{aligned}$$

now we get

$$|\widehat{\Lambda}_n - U_1| \leq Mh, \text{ where } M > 0 \text{ is a constant.}$$

By using a similar inequality like that of (13) with  $\widehat{\Lambda}_n, \Lambda$  and  $U_1$ , Lemma 1 and hypothesis  $H_1$  complete the last part of this proposition for  $n$  large enough.  $\blacktriangleleft$

*Proof. (Theorem 1)*

(i) Thanks to hypothesis  $A_2$ ,  $A_5 - i$ , we can write

$$\begin{aligned} |\widehat{A}_n(\theta) - A(\theta)| &= \left| \left[ \frac{\widehat{\Delta}_n}{\widehat{\mu}_n^{1-\theta}} \right]^{\frac{1}{1-\theta}} - \left[ \frac{\Delta}{\mu^{1-\theta}} \right]^{\frac{1}{1-\theta}} \right| \\ &\leq \left( \frac{1}{1-\theta} \right)^\kappa \left| \frac{\widehat{\Delta}_n}{\widehat{\mu}_n^{1-\theta}} - \frac{\Delta}{\mu^{1-\theta}} \right| \\ &\leq \left( \frac{1}{1-\theta} \right)^\kappa \left\{ \mu^{1-\theta} \left| \frac{1}{\widehat{\mu}_n^{1-\theta}} - \frac{1}{\mu^{1-\theta}} \right| + \frac{1}{\liminf_{n \leftarrow +\infty} \widehat{\mu}_n^{1-\theta}} |\widehat{\Delta}_n - \Delta| \right\} \\ &\leq M \left[ \left| \frac{1}{\widehat{\mu}_n^{1-\theta}} - \frac{1}{\mu^{1-\theta}} \right| + |\widehat{\Delta}_n - \Delta| \right], \end{aligned}$$

where

$$0 < M = \max \left\{ \mu^{1-\theta}; \frac{1}{\liminf_{n \leftarrow +\infty} \widehat{\mu}_n^{1-\theta}} \right\} \times \left( \frac{1}{1-\theta} \right)^\kappa.$$

Lemma 1 associated with the Mapping Theorem allows us to obtain

$$\left| \frac{1}{\widehat{\mu}_n^{1-\theta}} - \frac{1}{\mu^{1-\theta}} \right| \rightarrow 0, \text{ a.s. } n \leftarrow +\infty. \quad (14)$$

Thus (14) and Proposition 1 – (i) completes the first part of this theorem.

(ii) As previously in (i), the same assumptions give us:

$$\begin{aligned} |\widehat{A}(1) - A(1)| &= \left| \frac{\widehat{\Lambda}_n}{\widehat{\mu}} - \frac{\Lambda}{\mu} \right|, \\ &\leq \mu \left| \frac{1}{\widehat{\mu}} - \frac{1}{\mu} \right| + \frac{1}{\liminf_{n \leftarrow +\infty} \widehat{\mu}_n} |\widehat{\Lambda}_n - \Lambda| \\ &\leq M \left\{ \left| \frac{1}{\widehat{\mu}} - \frac{1}{\mu} \right| + |\widehat{\Lambda}_n - \Lambda| \right\}, \end{aligned}$$

where

$$0 < M = \left\{ \mu; \frac{1}{\liminf_{n \leftarrow +\infty} \widehat{\mu}_n} \right\}.$$

As previously in proof (i), Lemma 1 associated with the mapping theorem and Proposition 1 – (ii) complete the last part of this theorem.  $\blacktriangleleft$

*Proof. (Corollary 1)*

- (i) This first part will concern the case where  $\theta \in [0; 1[$ . Lemma 2 and the Triangle Inequality lead us to

$$\begin{aligned} |\widehat{W}_n(\theta) - W(\theta)| &= \left| \left[ \widehat{\mu}_n(1 - \widehat{A}_n(\theta)) \right]^{1-\theta} - [\mu(1 - A(\theta))]^{1-\theta} \right| \\ &\leq (1 - \theta) \left| \widehat{\mu}_n(1 - \widehat{A}_n(\theta)) - \mu(1 - A(\theta)) \right|^{1-\theta} \\ &\leq (1 - \theta) \left\{ (1 + A(\theta)) |\widehat{\mu}_n - \mu| + \widehat{\mu}_n |\widehat{A}_n(\theta) - A(\theta)| \right\} \\ &\leq M_\theta \left\{ |\widehat{\mu}_n - \mu| + |\widehat{A}_n(\theta) - A(\theta)| \right\}, \end{aligned}$$

where

$$0 < M_\theta = \max \left\{ 1 + A(\theta); \limsup_{n \leftarrow +\infty} \widehat{\mu}_n \right\} \times (1 - \theta), \quad \forall \theta \in [0; 1[.$$

Thus Lemma 1 and Theorem 1 – (i) complete the first part of this corollary.

- (ii) This second and last part concerns the case where  $\theta = 1$ . For this case especially, we assume that  $A(1)$  and its estimator  $\widehat{A}_n(1)$  are all different from 1 or all do not approach 1. Therefore  $\mu(1 - A(1))$  and  $\widehat{\mu}_n(1 - \widehat{A}_n(1))$  are strictly positive. Let us denote  $\delta = \min \left\{ \mu(1 - A(1)); \widehat{\mu}_n(1 - \widehat{A}_n(1)) \right\}$ . We therefore have  $\delta$  strictly positive. Thus the function  $z \mapsto \log(z)$  is  $\frac{1}{\delta}$ -Lipschitzian on  $[\delta; +\infty)$  because its derivative in absolute value is bounded above by  $\frac{1}{\delta}$ . This leads us to write:

$$\begin{aligned} |\widehat{W}_n(1) - W(1)| &= \left| \log[\widehat{\mu}_n(1 - \widehat{A}_n(1))] - \log[\mu(1 - A(1))] \right| \\ &\leq \frac{1}{\delta} \left| \widehat{\mu}_n(1 - \widehat{A}_n(1)) - \mu(1 - A(1)) \right| \\ &\leq M_\delta \left\{ |\widehat{\mu}_n - \mu| + |\widehat{A}_n(1) - A(1)| \right\}, \end{aligned}$$

where

$$0 < M_\delta = \frac{\max \left\{ 1 + A(1); \limsup_{n \leftarrow +\infty} \widehat{\mu}_n \right\}}{\delta}.$$

Thus Lemma 1 and Theorem 1 – (ii) complete the last part of this corollary.  $\blacktriangleleft$

*Proof. (Corollary 2)*

- For  $\theta \in [0, 1[$ , we directly and successively draw these inequalities from the following relation (see proof of the Proposition 1):

$$\widehat{\Delta}_n(\theta) \leq Q_{1-\theta} \left[ \frac{1}{n} \sum_{i=1}^n X_i^{1-\theta} + h^{1-\theta} \int_{\mathbb{R}} w^{1-\theta} K(w) dw \right].$$

Using the property of monotonic increasing and the linearity of the mathematical expectation and also  $||a| - |b|| \leq |a - b|$ , then we have

$$\begin{aligned} \mathbb{E} \left[ \widehat{\Delta}_n(\theta) \right] &\leq Q_{1-\theta} \mathbb{E} \left[ X^{1-\theta} \right] + h^{1-\theta} Q_{1-\theta} \nu(K) \\ &\Rightarrow \mathbb{E} \left| \widehat{\Delta}_n(\theta) - \Delta(\theta) \right| = \mathcal{O} \left( h^{1-\theta} \right). \end{aligned} \quad (15)$$

Moreover, we have

$$\begin{aligned} |\widehat{A}_n(\theta) - A(\theta)| &\leq M \left[ \left| \frac{1}{\widehat{\mu}_n} - \frac{1}{\mu} \right| + \left| \widehat{\Delta}_n(\theta) - \Delta(\theta) \right| \right] \\ \Rightarrow \mathbb{E} |\widehat{A}_n(\theta) - A(\theta)| &\leq M \left[ \mathbb{E} \left| \frac{1}{\widehat{\mu}_n} - \frac{1}{\mu} \right| + \mathbb{E} \left| \widehat{\Delta}_n(\theta) - \Delta(\theta) \right| \right]. \end{aligned}$$

Thus for  $n$  large enough, we have  $\mathbb{E} |\widehat{A}_n(\theta) - A(\theta)| \rightarrow 0$ . In particular we obtain

$$\mathbb{E} \left[ \widehat{A}_n(\theta) - A(\theta) \right] \rightarrow 0 \iff \mathbb{E} \left[ \widehat{A}_n(\theta) \right] \rightarrow \mathbb{E} [A(\theta)] = A(\theta).$$

We have just shown that the estimator is asymptotically unbiased. We can notice that  $(a - b)^2 \leq a^2 + b^2$  (where  $a$  and  $b$  have the same sign) and also the function  $z \mapsto z^p$ ,  $p \geq 2$  is convex (this allows us to apply Jensen's inequality). All this allows us to write

$$\begin{aligned} \left( \widehat{A}_n(\theta) - A(\theta) \right)^2 &\leq \left[ \int_{\mathbb{R}} \left( \frac{x}{\widehat{\mu}_n} \right)^{1-\theta} \widehat{f}_n(x) dx \right]^{\frac{2}{1-\theta}} + \left[ \int_{\mathbb{R}} \left( \frac{x}{\mu} \right)^{1-\theta} f(x) dx \right]^{\frac{2}{1-\theta}} \\ &\leq \int_{\mathbb{R}} \left( \frac{x}{\widehat{\mu}_n} \right)^2 \widehat{f}_n(x) dx + \int_{\mathbb{R}} \left( \frac{x}{\mu} \right)^2 f(x) dx \\ &\leq \frac{1}{\widehat{\mu}_n^2} \left[ \frac{1}{n} \sum_{i=1}^n X_i^2 \int_{\mathbb{R}} K(w) dw + 2h \frac{1}{n} \sum_{i=1}^n X_i^2 \int_{\mathbb{R}} wK(w) dw + h^2 \int_{\mathbb{R}} w^2 K(w) dw \right] + \frac{\mu_2}{\mu^2} \\ &\Rightarrow \mathbb{E} \left[ \left( \widehat{A}_n(\theta) - A(\theta) \right)^2 \right] = \mu_2 + \frac{\mu_2}{\mu^2} + \mathcal{O} \left( h^2 \right) = (\mu^2 + 1) \left( 1 + \frac{\sigma^2}{\mu^2} \right) + \mathcal{O} \left( h^2 \right). \end{aligned}$$

Furthermore we know that

$$\text{Var} \left[ \widehat{A}_n(\theta) - A(\theta) \right] = \mathbb{E} \left[ \left( \widehat{A}_n(\theta) - A(\theta) \right)^2 \right] - \mathbb{E}^2 \left[ \widehat{A}_n(\theta) - A(\theta) \right].$$

So, from the relation (15) we have

$$\text{Var} \left[ \widehat{A}_n(\theta) - A(\theta) \right] = (\mu^2 + 1) \left( 1 + \frac{\sigma^2}{\mu^2} \right) + \mathcal{O} \left( h^{2-2\theta} \right) \text{ a.s.}$$

For  $n$  large enough, we have  $\text{Var} \left[ \widehat{A}_n(\theta) - A(\theta) \right] \rightarrow s^2 = (\mu^2 + 1) \left( 1 + \frac{\sigma^2}{\mu^2} \right)$ .

◦ For  $\theta = 1$ , we also have (see proof of Proposition 1):

$$\widehat{\Lambda}_n \leq \frac{1}{n} \sum_{i=1}^n X_i + h \int_{\mathbb{R}} wK(w)dw, \Rightarrow \mathbb{E}[\widehat{\Lambda}_n] \leq \mathbb{E}[X_i] + h\nu(K)$$

$$\mathbb{E}|\widehat{\Lambda}_n - \mu| = \mathcal{O}(h) \text{ a.s.}$$

On the one hand the convexity of the exponential function, the triangular inequality and  $||a| - |b|| \leq |a - b|$  allows us to have

$$\mathbb{E}|\widehat{\Lambda}_n - \Lambda| \leq \mathbb{E}|\widehat{\Lambda}_n - \mu| + \mathbb{E}|\Lambda - \mu| \Rightarrow \mathbb{E}|\widehat{\Lambda}_n - \Lambda| = \mathcal{O}(h) \text{ a.s.}$$

On the other hand

$$|\widehat{\Lambda}_n(1) - A(1)| \leq M \left\{ \left| \frac{1}{\widehat{\mu}} - \frac{1}{\mu} \right| + |\widehat{\Lambda}_n - \Lambda| \right\}.$$

Thus, for  $n$  large enough

$$\mathbb{E}|\widehat{\Lambda}_n(1) - A(1)| = \mathcal{O}(h) \text{ a.s.} \quad (16)$$

We have also

$$\begin{aligned} \left( \widehat{A}_n(\theta) - A(\theta) \right)^2 &\leq \left\{ \frac{1}{\widehat{\mu}_n} \exp \left[ \int_{\mathbb{R}} \log(x) \widehat{f}_n(x) dx \right] \right\}^2 + \left\{ \frac{1}{\mu} \exp \left[ \int_{\mathbb{R}} \log(x) f(x) dx \right] \right\}^2 \\ &\leq \frac{1}{\widehat{\mu}_n^2} \exp \left[ 2 \int_{\mathbb{R}} \log(x) \widehat{f}_n(x) dx \right] + \frac{1}{\mu^2} \exp \left[ 2 \int_{\mathbb{R}} \log(x) f(x) dx \right] \\ &\leq \frac{1}{\widehat{\mu}_n^2} \int_{\mathbb{R}} x^2 \widehat{f}_n(x) dx + \frac{\mu_2}{\mu^2} \Rightarrow \mathbb{E} \left[ \left( \widehat{A}_n(\theta) - A(\theta) \right)^2 \right] \leq M [\mu_2 + h^2 \nu(K)] + \frac{\mu_2}{\mu^2}. \end{aligned}$$

For  $n$  large enough, we get

$$\Rightarrow \mathbb{E} \left[ \left( \widehat{A}_n(\theta) - A(\theta) \right)^2 \right] = \mu_2 + \frac{\mu_2}{\mu^2} + \mathcal{O}(h^2) \text{ a.s.} \quad (17)$$

Formulas (16) and (17) give the variance as previously  $\text{Var} \left[ \widehat{A}_n(\theta) - A(\theta) \right] = (1 + \mu^2) \left( 1 + \frac{\sigma^2}{\mu^2} \right) + \mathcal{O}(h^2) \text{ a.s.}$  This which completes the proof of this corollary. ◀

## 4. Simulation

Before moving on to numericals, we were able to calculate the Atkinson inequality index and welfare for certain probability distributions that will be useful for the rest.

Name	p.d.f. $f(x)$	Atkinson index $A(\theta)$	Welfare function $W(\theta)$
Beta $\mathcal{B}(\alpha, \beta)$	$\frac{1}{B(\alpha, \beta)} x^{\alpha-1} (1-x)^{\beta-1} 1_{]0,1[}(x)$	$\left\{ \begin{array}{l} 1 - \frac{\alpha + \beta}{\alpha} \left[ \frac{\Gamma(\alpha + 1 - \theta) \Gamma(\beta)}{B(\alpha, \beta) \Gamma(\alpha + \beta + 1 - \theta)} \right]^{\frac{1}{1-\theta}} \\ 1 - \frac{\alpha + \beta}{\alpha} \exp[\psi(a) - \psi(a + b)] \end{array} \right.$	$\left\{ \begin{array}{l} \frac{\Gamma(\alpha + 1 - \theta) \Gamma(\beta)}{B(\alpha, \beta) \Gamma(\alpha + \beta + 1 - \theta)} \\ \psi(a) - \psi(a + b) \end{array} \right.$
Exponential $\mathcal{E}(\lambda)$	$\lambda \exp(-\lambda x) 1_{]0,+\infty[}(x)$	$\left\{ \begin{array}{l} 1 - \Gamma^{\frac{1}{1-\theta}}(2 - \theta) \\ 1 - \lambda \exp[-(\gamma + \log(\lambda))] \end{array} \right.$	$\left\{ \begin{array}{l} \frac{\Gamma(2 - \theta)}{\lambda^{1-\theta}} \\ -(\gamma + \log(\lambda)) \end{array} \right.$
Gamma $\Gamma(\alpha, \beta)$	$\frac{\beta^\alpha}{\Gamma(\alpha)} x^{\alpha-1} \exp(-\beta x) 1_{]0,+\infty[}(x)$	$\left\{ \begin{array}{l} 1 - \frac{\Gamma^{\frac{1}{1-\theta}}(1 - \theta + \alpha)}{\alpha \Gamma^{\frac{1}{1-\theta}}(\alpha)} \\ 1 - \frac{\beta}{\alpha} \exp[\psi(\alpha) - \log(\beta)] \end{array} \right.$	$\left\{ \begin{array}{l} \frac{\Gamma(1 - \theta + \alpha)}{\beta^{1-\theta} \Gamma(\alpha)} \\ \psi(\alpha) - \log(\beta) \end{array} \right.$
Pareto $\mathcal{P}(\alpha, \beta)$	$\frac{\alpha \beta^\alpha}{x^{\alpha+1}} 1_{]0,+\infty[}(x)$	$\left\{ \begin{array}{l} 1 - \frac{\alpha - 1}{(\alpha + \theta - 1)^{\frac{1}{1-\theta}}} \\ 1 - \frac{\alpha - 1}{\alpha \beta} \exp \left[ \frac{\alpha \log(\beta) + 1}{\alpha} \right] \end{array} \right.$	$\left\{ \begin{array}{l} \frac{(\alpha \beta)^{1-\theta}}{\alpha + \theta - 1} \\ \frac{\alpha \log(\beta) + 1}{\alpha} \end{array} \right.$
Uniform $\mathcal{U}([\alpha, \beta])$	$\frac{1}{\beta - \alpha} 1_{]0,+\infty[}(x)$	$\left\{ \begin{array}{l} 1 - \frac{2}{\alpha + \beta} \left[ \frac{\beta^{2-\theta} - \alpha^{2-\theta}}{(\beta - \alpha)(2 - \theta)} \right]^{\frac{1}{1-\theta}} \\ 1 - \frac{2}{\alpha + \beta} \exp \left[ \frac{\beta \log(\beta) - \alpha \log(\alpha)}{\beta - \alpha} - 1 \right] \end{array} \right.$	$\left\{ \begin{array}{l} \frac{\beta^{2-\theta} - \alpha^{2-\theta}}{(\beta - \alpha)(2 - \theta)} \\ \frac{\beta \log(\beta) - \alpha \log(\alpha)}{\beta - \alpha} - 1 \end{array} \right.$
Weibull $\mathcal{W}(\alpha, \beta)$	$\frac{\alpha}{\beta} x^{\alpha-1} \exp\left(-\frac{x^\alpha}{\beta}\right) 1_{]0,+\infty[}(x)$	$\left\{ \begin{array}{l} 1 - \frac{\Gamma^{\frac{1}{1-\theta}}\left(1 + \frac{1-\theta}{\alpha}\right)}{\Gamma\left(1 + \frac{1}{\alpha}\right)} \\ 1 - \frac{1}{\beta^{\frac{1}{\alpha}}} \exp\left[-\frac{\log(\beta) + \gamma}{\alpha}\right] \end{array} \right.$	$\left\{ \begin{array}{l} \beta^{\frac{1-\theta}{\alpha}} \Gamma\left(1 + \frac{1-\theta}{\alpha}\right) \\ -\frac{\log(\beta) + \gamma}{\alpha} \end{array} \right.$

**Tab. 1.** Theoretical expressions of the Atkinson index and the welfare function

4. 1. Atkinson index for some usual probability distributions

However, theoretical calculations of the Atkinson index with probability distributions are not easy or easy to determine. With other distributions, this seems very complex or even impossible. We will spare you these details and the results obtained are grouped in the Table 1. The theoretical calculation of the Atkinson index involves certain so-called special functions which are defined by:

- Gamma function:  $\Gamma(a) = \int_0^\infty z^{a-1} \exp(-z) dz, a \in \mathbb{R}_+$
- Digamma function:  $\psi(a) = \int_0^\infty z^{a-1} \log(z) \exp(-z) dz = \int_0^\infty \left( \frac{e^{-z}}{z} - \frac{e^{-az}}{1 - e^{-z}} \right) dz = \frac{d}{da} \log[\Gamma(a)], a \in \mathbb{R}_+$
- Euler - Mascheroni constant:  $\gamma = -\psi(1) = -\int_0^\infty \log(z) e^{-z} dz = \lim_{n \rightarrow +\infty} \left( \sum_{k=1}^n \frac{1}{k} - \log(n) \right) \simeq 0.577216664901532860$



◦ Beta function:  $B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}$ ,  $a, b \in \mathbb{R}_+$

**Remark 2.** Recall that the first lines of the expressions that are in the braces represent the general case for  $\theta \neq 1$  and those of the second lines represent the particular case for  $\theta = 1$ .

The table being obtained, we choose one of these models to study the convergence of our nonparametric estimator with respect to that. Our choice is the Weibull model because of its particularity in terms of application and its multitasking. The Weibull distribution intervenes in almost all areas like that of Pareto in the estimations of income inequality measures (see Agbokou [5]). To do this, with the data we have, we will determine these parameters by the Maximum Likelihood (M.L.) method to obtain the theoretical Atkinson index.

#### 4. 2. Determination of the parameters of the Weibull distribution by the M.L. method

Due to its adaptability in fitting distributions or data particularly in data science, the Weibull distribution has taken in recent years a major position in the field of parametric, semi or nonparametric estimations. One of the main barriers to a more important or wider use of the Weibull distribution is the complexity of estimating its parameters. Regrettably or sadly, the calculations that this estimation involves are not always simple enough. This subsection deals with the estimation of maximum likelihood in samples of the Weibull probability density. The likelihood function of this sample  $(X_1, \dots, X_n)$  is given by:

$$L(X_1, \dots, X_n, (\alpha, \beta)) = \prod_{i=1}^n \frac{\alpha}{\beta} X_i^{\alpha-1} \exp\left(-\frac{X_i^\alpha}{\beta}\right).$$

We let  $\mathcal{L}(\alpha, \beta) = \mathcal{L}(X_1, \dots, X_n, (\alpha, \beta)) = \log L(X_1, \dots, X_n, (\alpha, \beta))$  represent the log-likelihood, then we derive  $\mathcal{L}(\alpha, \beta)$  respectively with respect to  $\alpha$  and with respect to  $\beta$  then by setting the two partial derivatives equal to zero each, we obtain:

$$\begin{cases} \frac{\partial \mathcal{L}(\alpha, \beta)}{\partial \alpha} = \frac{n}{\alpha} + \sum_{i=1}^n \log(X_i) - \frac{1}{\beta} \sum_{i=1}^n X_i^\alpha \log(X_i) = 0, \\ \frac{\partial \mathcal{L}(\alpha, \beta)}{\partial \beta} = -\frac{n}{\beta} + \frac{1}{\beta^2} \sum_{i=1}^n X_i^\alpha = 0. \end{cases} \quad (18)$$

On eliminating  $\beta$  between these two equations of the system (18) and simplifying the term, we get:

$$\frac{\sum_{i=1}^n X_i^\alpha \log(X_i)}{\sum_{i=1}^n X_i^\alpha} - \frac{1}{\alpha} - \frac{1}{n} \sum_{i=1}^n \log(X_i) = 0. \quad (19)$$

We note that equation (19) is difficult or even impossible to solve analytically. We will therefore use a numerical resolution or iterative methods such as the Newton method or the fixed point method or the secant and regula falsi method in order to find the estimator  $\hat{\alpha}$  of the parameter  $\alpha$ . If the estimator  $\hat{\alpha}$  is obtained, we therefore deduce from (18), the estimator  $\hat{\beta}$  of  $\beta$  by the equality:

$$\hat{\beta} = \frac{1}{n} \sum_{i=1}^n X_i^{\hat{\alpha}}.$$

#### 4. 3. Real data description and simulation results

For our study, we focused on the World Bank data that could be easily found on their website DataBank We were interested in the "adjusted net national income per capita (current US dollars)" of 2021 for each country in the world (at least for those who have it because we could not find the updated data for all countries). Since the data is allocated by region according to the World Bank's breakdown, we organized this data by continent as follows:

- Europe Zone (1) = European Union
- America Zone (2) = America and the Caribbean + North America
- Asia Zone (3) = East Asia and the Pacific + South Asia + Middle East + Central Asia
- Africa Zone (4) = Sub-Saharan Africa + North Africa
- Eurasia Zone (5) = Europe Zone + Asia Zone
- World (6) = the sum of all the above zones

**Remark 3.** The last two groupings aim not only to study the inequalities of countries between two countries or between countries of the world but also to see the performance of our estimators when the sample size increases and also the impact of the inversion parameter on them depending on these sizes.

Table 2 provides a statistical description of some position and dispersion characteristics of these data taken to the thousandth in order to facilitate the calculations of the programming in the rest of this subsection using Matlab software.

Statistic summary	Size	Minimum	Mean	Median	Mode	Maximum	Coefficient of variation
EUROPEAN AREA	27	7.6500	25.3004	19.6460	7.6500	65.3640	0.5857
AMERICA AREA	33	1.2510	10.6735	6.6080	1.2510	52.6190	0.9870
ASIA AREA	51	0.3400	9.7364	3.5040	0.3400	44.5130	0.7573
AFRICA AREA	54	0.1560	1.9479	1.1455	0.1560	12.2520	0.7102
EURASIAN AREA	78	0.3400	15.1239	9.0055	0.3400	65.3640	0.9823
WORLD	165	0.1560	9.9217	4.0190	0.4860	65.3640	0.7618

**Tab. 2.** *Statistical description of data from the World Bank website*

In view of the statistical summary of Table 2, we notice that the data, whatever the area, are very heterogeneous since the coefficient of variation of each of these groupings is strictly greater than 0.30. This heterogeneity character makes this study very exciting

since with simulated data (which we do not present here), we always had homogeneous data which contribute to the robustness of our estimators from the size  $n = 30$ . Thus the other objective is to see the behavior of our estimators facing heterogeneous data because we know that heterogeneous data in statistical study such as estimates and forecasts negatively affect or slow down the robustness of the estimators and by ricochet impact the forecasts.

Thus, for the simulation of our estimators, we chose as a statistical kernel  $K$ , the Epanechnikov kernel also called the "parabolic" kernel. It is named after the author Epanechnikov [12], who used and studied it for the first time in 1969. On the one hand, it is well known in the literature that the kernel has little influence on the performance of nonparametric estimators. What motivates this choice is that this kernel allows to have the most efficient estimator for the density, which concerns us in this document. On the other hand, the literature reveals that the choice of the smoothing parameter  $h = h_n$  has a major influence on the robustness of the kernel estimator (see Agbokou et al. [2], [3]). The process of this choice in the presence of real data is not as easy as in the presence of simulated data. We therefore opted for the numerical cross-validation method which consists in minimizing the integrated squared error defined from a series of observations or data  $(x_i)_{1 \leq i \leq n}$  of size  $n$ . This error is defined by:

$$\Phi(h_n) = \int_{x_{(1)}}^{x_{(n)}} \left( \widehat{A}_n(\theta) - A(\theta) \right)^2 dx,$$

where  $x_{(1)} = \min(x_i)$  and  $x_{(n)} = \max(x_i)$  for all  $i \in \{1, \dots, n\}$ .

The bandwidth parameter selection rule results in the minimization of this criterion:

$$\widehat{h}_n = \arg \min_{h_n} \Phi(h_n). \quad (20)$$

The smoothing parameter of (20) thus obtained is asymptotically optimal. To determine the parameter  $h$ , our Matlab code is programmed in such a way that we obtain at the same time the RMSE (Residual Mean Square Error). This is the root mean square deviation, which is the standard deviation of the residuals (prediction errors). The residuals are the measure of the deviation between the data points and the regression line. The RMSE metric is the measure of the distribution of these residuals. In other words, it indicates the concentration of the data around the line of best fit. In our work we have two parameters  $h$  to determine and therefore two RMSEs. We denote them respectively  $h_{1n}$  for the estimator of the Atkinson index and  $h_{2n}$  for its associated welfare function. Similarly, their errors or residuals are respectively noted  $RMSE_1$  and  $RMSE_2$ . To get an idea of the impact of the aversion parameter, we chose the parameter  $\theta$  such that  $\theta \in \{0.01, 0.05, 0.1, 0.5, 0.9, 1\}$  for each sample size (or area or region). We also calculated for each  $\theta$  the classical estimators (usual classical forms) of the Atkinson index (6) and its welfare function (5) in order to see if they best estimate the theoretical Atkinson index and its theoretical welfare function.

Let us recall that the two parameters of the theoretical Atkinson index and its theoretical welfare function cannot be taken arbitrarily, we opted for the maximum likelihood (M.L.) method with the intervention of Newton's numerical method in order to be able to assign a value to each of these parameters.

The results obtained are summarized in Table 3. A visualization of all these results obtained is summarized in Figure 2.

$\theta$	$\hat{h}_{1n}$	$\hat{h}_{2n}$	$\hat{\alpha}$	$\hat{\beta}$	$A(\theta)$	$\hat{A}_n(\theta)$	$\bar{A}(\theta)$	$W(\theta)$	$\hat{W}_n(\theta)$	$\bar{W}(\theta)$	$RMSE_1$	$RMSE_2$
EUROPEAN AREA												
0.01	0.4609	0.4268	1.8719	535.6642	0.0016	0.0015	0.0972	24.5977	24.0057	22.1368	0.0092	0.0560
0.05	0.4701	0.4193	1.8719	535.6642	0.0078	0.0077	0.1035	21.3725	20.4040	22.4932	0.0081	3.8750
0.1	0.4609	0.4191	1.8719	535.6642	0.0157	0.0154	0.1115	17.9269	16.4668	20.0677	0.0094	2.1319
0.5	0.8231	0.5110	1.8719	535.6642	0.0815	0.0816	0.0760	4.3684	4.5351	9.6700	3.08e-08	0.0278
0.9	0.2676	0.0879	1.8719	535.6642	0.1534	0.1534	0.1337	1.0515	1.0516	3.6170	1.15e-11	1.77e-08
1	0.7500	0.5036	1.8719	535.6642	0.9991	0.9698	0.5432	3.6651	3.0713	3.0708	0.2078	0.3532
AMERICA AREA												
0.01	0.9000	0.0224	1.1726	17.2937	0.0033	0.0036	0.0037	10.4673	10.4673	10.4908	8.59e-08	2.13e-18
0.05	0.9000	0.1859	1.1726	17.2937	0.0166	0.0182	0.0185	9.3710	9.3174	9.8079	3.57e-06	0.0029
0.1	0.5255	0.2041	1.1726	17.2937	0.0332	0.0362	0.0364	8.1578	8.1482	9.0539	1.01e-05	9.21e-05
0.5	0.5290	0.4391	1.1726	17.2937	0.1703	0.1704	0.1682	2.6477	2.8142	5.9593	7.36e-10	0.0277
0.9	0.9116	0.0824	1.1726	17.2937	0.3160	0.3160	0.2799	0.8255	0.8261	2.2623	8.27e-12	3.33e-07
1	0.6641	0.1854	1.1726	17.2937	0.9953	0.4203	0.4205	2.9232	2.0131	2.0042	0.3304	0.8464
ASIA AREA												
0.01	0.6464	0.3758	0.7907	5.3641	0.0060	0.0060	0.0068	9.3052	9.4544	9.5485	4.36e-10	0.0223
0.05	0.2968	0.3760	0.7907	5.3641	0.0300	0.0300	0.0340	8.3408	8.4081	8.8504	1.53e-10	0.0045
0.1	0.8701	0.3802	0.7907	5.3641	0.0599	0.0598	0.0678	7.2691	7.2782	8.0883	2.84e-09	8.25e-05
0.5	0.3809	0.2296	0.7907	5.3641	0.2950	0.3200	0.3246	2.3305	2.4987	5.1461	8.78e-04	0.0283
0.9	0.3839	0.2891	0.7907	5.3641	0.5226	0.5176	0.5596	0.6747	0.8069	1.6730	0.0014	0.0175
1	0.0009	0.3833	0.7907	5.3641	0.9931	0.9837	0.3153	2.8545	2.8535	1.4622	8.95e-05	1.04e-06
AFRICA AREA												
0.01	0.7214	0.6738	1.0477	22.9593	0.0039	0.0034	0.0045	18.8885	18.8885	19.0143	2.91e-07	7.28e-18
0.05	0.6070	0.5713	1.0477	22.9593	0.0197	0.0226	0.1340	16.4931	17.2962	14.6465	0.0131	3.4100
0.1	0.5973	0.5712	1.0477	22.9593	0.0395	0.0447	0.1114	13.9155	12.4864	13.4340	0.0125	2.0423
0.5	0.7062	0.7059	1.0477	22.9593	0.2008	0.2068	0.2151	3.5000	3.7051	7.8615	2.04e-04	0.0421
0.9	0.5804	0.9853	1.0477	22.9593	0.3692	0.3395	0.4800	0.8351	0.8648	6.9105	0.0123	8.80e-04
1	0.7422	0.5658	1.0477	22.9593	0.9985	0.5211	0.3089	3.5419	2.5103	2.4167	0.0123	8.80e-04
EURASIAN AREA												
0.01	0.5486	0.9742	0.8894	10.6895	0.0050	0.0049	0.0047	14.7255	14.7255	14.8590	1.12e-07	2.50e-10
0.05	0.8626	0.3224	0.8894	10.6895	0.0252	0.0248	0.0242	12.9685	12.9685	13.6272	1.92e-06	9.99e-15
0.1	0.8565	0.0523	0.8894	10.6895	0.0505	0.0499	0.0496	11.0575	11.0575	12.2792	7.14e-07	1.98e-09
0.5	0.5156	0.9413	0.8894	10.6895	0.2524	0.2574	0.2615	2.9995	3.3016	6.7171	8.21e-05	0.0912
0.9	0.5163	0.5708	0.8894	10.6895	0.4554	0.4598	0.5088	0.7528	0.8441	2.3427	0.0028	0.0083
1	0.0001	0.8033	0.8894	10.6895	0.9975	0.9918	0.5044	3.3128	3.3128	2.0155	0.0028	0.0083
WORLD												
0.01	0.5662	0.4996	0.7660	5.0871	0.0063	0.0058	0.0068	9.5207	9.6375	9.7281	2.35e-07	0.0137
0.05	0.8813	0.5002	0.7660	5.0871	0.0314	0.0314	0.0340	8.5243	8.5644	9.0107	2.73e-09	0.0016
0.1	0.2698	0.8729	0.7660	5.0871	0.0627	0.0627	0.0678	7.4182	7.4182	8.2268	1.20e-11	3.56e-12
0.5	0.5095	0.5512	0.7660	5.0871	0.3072	0.3261	0.3347	2.3449	2.5396	5.1714	7.58e-04	0.0379
0.9	0.4223	0.3933	0.7660	5.0871	0.5411	0.5401	0.6442	0.6642	0.8083	0.6392	0.0106	0.0208
1	0.0029	0.5095	0.7660	5.0871	0.9933	0.9841	0.1288	2.8771	2.8771	1.4180	8.49e-05	1.02e-11

Tab. 3. Comparison of theoretical values and estimates of the Atkinson index and the welfare





Fig. 2. Comparison of theoretical and estimates curves of the Atkinson index and the welfare

#### 4. 4. Discussion

First of all, we note that kernel estimators provide very good adjustments for large samples, i.e. from size  $n = 70$  and these provide better adjustments compared to those that are classical. Apparently these estimators are less sensitive (evolve linearly) for values less than 1 of the aversion parameters compared to those that are classical. In addition, we note that, the more the parameter increases, the more the indices increase sharply and their welfare functions decrease slightly. For  $\theta = 1$  all the estimators, even those qualified as classical, are very sensitive and their values sometimes differ in terms of the direction of variations (some increase when the others decrease for the Atkinson indices). Generally speaking, for Theta close to 0, the values of the estimators are very small and in the neighborhood of 1 or at 1, they are very large. This character shows the major impact of the aversion parameter for the study of inequalities in a population and it poses a great debate because a bad choice of the aversion parameter can lead us to make a hasty conclusion or one that may be far from reality. Thus for a global view, we calculated the average of all the values of the Atkinson index, which is provided by the following Figure 3:



**Fig. 3. Diagram of the average Atkinson indices and average welfare values**

This last figure shows that the Atkinson index whatever the type of estimator does not exceed 0.35 in each zone and therefore on each continent. Although these averages do not allow us to conclude effectively or to draw an efficient objective conclusion, nevertheless they allow us to have more or less or approximately an overall view of the trend of the Atkinson indices and the values of the associated welfare in each region or zone. On the one hand, we therefore conclude that Figure 2 and Table 3 have provided us with the behavior of our estimators in relation to the progressive choice of the aversion parameter

$\theta$ . On the other hand, to estimate the Atkinson indices and the values of social welfare, it is therefore essential to find an estimator of the parameter  $\theta$  or therefore find the corresponding value of  $\theta$  for each data or observation. Thus we therefore propose here to find the value of  $\theta$  for each of our observations. To do this, the function  $\Phi$  (4. 3.) therefore becomes a bivariate function whose variables are the bandwidth and aversion parameters. The optimization problem (20) is therefore a bivariate optimization problem on the unit square surface and becomes:

$$\left(\hat{h}_n, \hat{\theta}_n\right) = \arg \min_{h_n, \theta \in [0,1]} \Phi(h_n, \theta).$$

Let  $\hat{\theta}_{1n}$  and  $\hat{\theta}_{2n}$  be the respective aversion parameters of the Atkinson index and welfare. The only parameters that do not change compared to the previous results are the parameters  $\hat{\alpha}$  and  $\hat{\beta}$  from the M.L method. The results obtained are grouped in Table 4, giving the values of the Atkinson indices and those of welfare after estimating the aversion parameters. Since the classical Atkinson indices and associated welfare function do not provide a very good fit, we no longer present it in this table.

Area or Zone	$\hat{h}_{1n}$	$\hat{\theta}_{1n}$	$RMSE_1$	$\hat{A}_n(\theta)$	$A(\theta)$	$\hat{h}_{2n}$	$\hat{\theta}_{2n}$	$RMSE_2$	$\hat{W}_n(\theta)$	$W(\theta)$
EUROPE	0.4495	1.0000	0.0127	<b>0.2849</b>	0.2981	0.9604	0.0851	0.0759	<b>2.8102</b>	2.8233
AMERICA	0.6547	0.5329	1.79e-10	<b>0.1820</b>	0.1820	0.8718	0.1152	0.0002	<b>7.8195</b>	7.8195
ASIA	0.5710	0.7049	7.42e-08	<b>0.4125</b>	0.4125	0.0458	0.1140	2.13e-09	<b>0.9093</b>	0.9093
AFRICA	0.3894	0.6457	0.0018	<b>0.2606</b>	0.2612	0.5649	0.4643	0.0033	<b>3.9650</b>	3.9653
EURASIA	0.5571	0.3962	2.33e-11	<b>0.2034</b>	0.2034	0.4721	0.3316	9.07e-07	<b>5.2194</b>	5.2194
WORLD	0.5306	0.4209	1.05e-07	<b>0.2604</b>	0.2604	0.7792	0.0939	6.56e-10	<b>7.5482</b>	7.5482

**Tab. 4.** *Atkinson indices and welfare values based on the estimated aversion parameter*

Table 4 allows us to draw a more or less effective conclusion because it corresponds to the true trends of the Atkinson index. However, we can see that in each zone the aversion parameter is around 0.5 on average for the Atkinson index except for the Europe zone. This could be explained by the fact that the sample size is small, i.e. less than 50. On the other hand, for large samples, it is lower. As for welfare, the aversion parameter is on average 0.3. This diversity of values taken by the aversion parameter shows once again the importance of its estimation because it has more influence on the estimators. Thus, compared to the previous results (where  $\theta$  is taken arbitrarily), we see that there is a slight significant difference because the indices in this case are around 0.35 or even exceed it. However, it can be noted that inequality is more glaring in the Americas than on other continents. This may be due to the per capita income of North America which is more than ten times higher compared to those of the countries of the South or the Caribbean. In Africa and Europe, this inequality is almost identical to that in the world. Inequality is less reduced in Asia and this could be explained by the almost identical per capita income of several countries of the Gulf and the Pacific. Overall, the disparity is significant on all continents and world leaders must make much more efforts to raise the level of net national income per capita so that the inequality index on each continent or even in each country is at least higher than 0.7 by taking appropriate decisions for the development of each country.

## 5. Conclusion

In this paper, we have done a little more in-depth study on the Atkinson index and its welfare function than we had thought at the beginning of this study. First, we focused on the construction of the two estimators and then the study of their almost spure asymptotic convergence as well as asymptotic normality. Then, we looked at the simulation study which confirms the robustness of the two estimators for large sample sizes by working with real data which are very heterogeneous. These studies revealed that the aversion parameter has a huge influence on the estimation of the Atkinson index and cannot be chosen randomly or arbitrarily for an efficient study. We have therefore proposed a numerical estimation of said parameter by the cross-validation method in order to overcome this problem of choosing the aversion parameter. Finally, our study has shown that disparity in the world is a current problem and that it is not to be remedied in the short term but it is a collective problem and to be resolved in the long term. As a consequence or open problem of this study, it should be noted that it is therefore important to find a way to analytically estimate the aversion parameter using either methods that already exist such as the maximum likelihood method, the method of moments or the Bayesian method (as in Agbokou and al. [4]) or to find others (as in Dabana and al. [11]) that could better help us, and this will be the subject of our future work.

**Acknowledgement.** The authors thank the reviewers for their insightful comments and excellent work.

## References

1. Agbokou K. On the nature of elasticity function: an investigation and a kernel estimation. *J. Appl. Math.*, 2023, **2023** (Art. ID 1346602), pp. 1-14.
2. Agbokou K., Gneyou K.E. On the strong convergence of the hazard rate and its maximum risk point estimators in presence of censorship and functional explanatory covariate. *Afr. Stat.*, 2017, **12** (3), pp. 1397-1416.
3. Agbokou K., Gneyou K.E., Deme El H. Almost sure representations of the conditional hazard function and its maximum estimation under right-censoring and left-truncation. *Far East J. Theor. Stat.*, 2018, **54** (2), pp. 141-173.
4. Agbokou K., Mensah Y. Inference on the reproducing kernel Hilbert spaces. *Universal J. Math. Math. Sci.*, 2022, 15, pp. 11-29.
5. Agbokou K., Mensah Y. Varying bandwidth parameter method on kernel Gini index estimation. *An. Univ. Craiova Ser. Mat. Inform.*, 2023, **50** (1), pp. 29-41.
6. Atkinson A.B. On the measurement of inequality. *J. Economic Theory*, 1970, **2** (3), pp. 244-263.
7. Atkinson A.B. *The Economics of Inequality*. Oxford Univ. Press, London, 1975.
8. Banerjee A. A multidimensional Gini index. *Math. Social Sci.*, 2010, **60** (2), pp. 87-93.
9. Bellù L.G., Liberati P. *Policy Impacts on Inequality: Welfare Based Measures of Inequality - The Atkinson Index*. EASYPol, Module 050, FAO, 2006.



10. Biewen M., Jenkins S. *Estimation of Generalized Entropy and Atkinson Inequality Indices from Complex Survey Data*, No. 345, Discussion Papers of DIW Berlin, DIW Berlin, German Institute for Economic Research, 2003.
11. Dabana H., Agbokou K., Gneyou K.E. Kernel estimation of the conditional probability density function and mode in presence of functional covariate and censorship. *Afr. Stat.*, 2024, **19** (1), pp. 3771-3796.
12. Epanechnikov V.A. Non-parametric estimation of a multivariate probability density. *Theory Probab. Appl.*, 1969, **14** (1), pp. 153-158.
13. Guerrero V.M. A note on the estimation of Atkinson's index of inequality. *Economics Letters*, 1987, **25**, pp. 379-384.
14. Harper G., Price R. *A Framework for Understanding the Social Impacts of Policy and Their Effects on Wellbeing - A Paper for the Social Impact Taskforce*. Defra Evidence and Analysis Series, Paper 3, Defra, London, 2011.
15. Lehmann E.L. *Elements of Large-Sample Theory*. Springer, New York, 1998.
16. Piketty T. *Les Hauts Revenus en France au XX<sup>e</sup> Siècle: Inégalités et Redistributions, 1901-1998*. Grasset, Paris, 2001 (in French).
17. Rawls J. *A Theory of Justice*. Oxford Univ. Press, London, 1972.
18. Sen A. *On Economic Inequality*. Clarendon Press, Oxford, 1973.
19. Sen A. *On Economic Inequality*. Clarendon Press, Oxford, 2nd edition, 1997.
20. Serfling R.J. *Approximation Theorems of Mathematical Statistics*. The Johns Hopkins University, John Wiley & Sons, Inc., 1998.
21. Tchamyu V.S. Education, lifelong learning, inequality and financial access: evidence from African countries. *Contemp. Social Sci.*, 2020, **15** (1), pp. 7-25.