

## THE BOUNDEDNESS OF MAXIMAL OPERATORS IN CALDERON WEIGHTED $B$ -MORREY SPACES

S.H. KASUMOVA, S.Z. KHALIGOVA\*

Received: 18.09.2025 / Revised: 17.11.2025 / Accepted: 24.11.2025

**Abstract.** *In this paper the boundedness of the  $B$ -maximal operator  $M_\gamma$  from the  $B$ -Morrey space  $L_{p,\lambda,\omega,\gamma}(\mathbb{R}_{k,+}^n)$  to  $L_{p,\lambda,\omega,\gamma}(\mathbb{R}_{k,+}^n)$  for all  $1 < p < \infty$  when  $\omega \in C_{p,\gamma}(\mathbb{R}_{k,+}^n)$  is proved.*

**Keywords:**  $B$ -maximal operator, Laplace-Bessel differential operator, generalized shift operator, weighted Lebesgue space, Calderon weighted Morrey space

**Mathematics Subject Classification (2020):** 42B20, 42B25, 42B35

### 1. Introduction

The maximal operator, potentials, singular integrals and related topics associated with the Laplace-Bessel differential operator

$$\Delta_B = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} + \sum_{i=1}^k \frac{\gamma_i}{x_i} \frac{\partial}{\partial x_i}, \quad \gamma_1 > 0, \dots, \gamma_k > 0,$$

have been investigated by many researchers, see B. Muckenhoupt and E. Stein [21], I. Kipriyanov [17], K. Triméche [25], L. Lyakhov [19], K. Stempak [24], A. D. Gadjiev and I. A. Aliev [8], I. A. Aliev and S. Bayrakci [2], V. S. Guliyev [10], [11], V. S. Guliyev and J. J. Hasanov [12], [13], J. J. Hasanov and Z. V. Safarov [15], A. Şerbetçi and I. Ekincioglu [22], E. L. Shishkina [23] and others.

$B$ -Riesz potentials  $I_{\alpha,\gamma}$  generated by the generalized shift operator play important tool in Fourier-Bessel harmonic analysis and applications. These spaces have been investigated by lots of mathematicians [1], [2]-[5], [10], [11], [14], [16], [22]. In the present article

---

\* Corresponding author.

**Sabina H. Kasumova**

Azerbaijan State Pedagogical University, Baku, Azerbaijan  
E-mail: sabine.kasumova.adpu@mail.ru

**Sevinc Z. Khaligova**

Azerbaijan State Pedagogical University, Baku, Azerbaijan  
E-mail: seva.xaligova@hotmail.com

we give the necessary and sufficient conditions for the boundedness of the commutators of the  $B$ -Riesz potentials,  $[b, I_{\alpha, \gamma}]$  where  $b$  is a locally integrable function on  $\mathbb{R}_{k,+}^n$ , from the spaces  $L_{p, \gamma}(\mathbb{R}_{k,+}^n)$  to  $L_{q, \gamma}(\mathbb{R}_{k,+}^n)$ ,  $1 < p < \frac{n+|\gamma|}{\alpha}$ ,  $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n+|\gamma|}$  and from the spaces  $L_{1, \gamma}(\mathbb{R}_{k,+}^n)$  to  $WL_{q, \gamma}(\mathbb{R}_{k,+}^n)$ ,  $1 - \frac{1}{q} = \frac{\alpha}{n+|\gamma|}$ . Also, It is proved that  $B$ -maximal commutators  $M_{b, \gamma}$  and commutators of  $B$ -singular integral operators  $[b, A_{\gamma}]$  are bounded from the  $B$ -Morrey space  $L_{p, \lambda, \gamma}(\mathbb{R}_{k,+}^n)$  to  $L_{p, \lambda, \gamma}(\mathbb{R}_{k,+}^n)$  for all  $1 < p < \infty$ ,  $b \in BMO_{\gamma}(\mathbb{R}_{k,+}^n)$ .

Morrey spaces,  $L_{p, \lambda}(\mathbb{R}^n)$  introduced by C. Morrey in 1938 [20] play an important role in the theory of partial differential equations. These are defined by the following norm,  $f \in L_p^{loc}(\mathbb{R}^n)$

$$\|f\|_{L_{p, \lambda}} \equiv \|f\|_{L_{p, \lambda}(\mathbb{R}^n)} = \sup_{x \in \mathbb{R}^n, r > 0} r^{-\frac{\lambda}{p}} \|f\|_{L_p(B(x, r))} < \infty,$$

where  $0 \leq \lambda \leq n$ ,  $1 \leq p < \infty$ . If  $\lambda = 0$  and  $\lambda = n$  then  $L_{p, \lambda}(\mathbb{R}^n) = L_p(\mathbb{R}^n)$  and  $L_{p, \lambda}(\mathbb{R}^n) = L_{\infty}(\mathbb{R}^n)$  respectively. If  $\lambda < 0$  or  $\lambda > n$  then  $L_{p, \lambda}(\mathbb{R}^n) = \Theta$  where  $\Theta$  is the set of all functions equivalent to 0 on  $\mathbb{R}^n$ .

These spaces appeared to be quite useful in the study of the local behaviour of the solutions to elliptic partial differential equations, apriori estimates and other topics in the theory of partial differential equations.

F. Chiarenza and M. Frasca [7] studied the boundedness of the maximal operator in Morrey spaces.

Also we denote by  $WL_{p, \lambda}(\mathbb{R}^n)$ , the weak Morrey space, is the space of all functions  $f \in WL_p^{loc}(\mathbb{R}^n)$  for which

$$\|f\|_{WL_{p, \lambda}} \equiv \|f\|_{WL_{p, \lambda}(\mathbb{R}^n)} = \sup_{x \in \mathbb{R}^n, r > 0} r^{-\frac{\lambda}{p}} \|f\|_{WL_p(B(x, r))} < \infty,$$

where  $WL_p(\mathbb{R}^n)$  denotes the weak  $L_p$ -space.

## 2. Preliminaries

Let  $\mathbb{R}_{k,+}^n = \{x = (x_1, \dots, x_n) \in \mathbb{R}^n : x_1 > 0, \dots, x_k > 0, 1 \leq k \leq n\}$  and let  $E(x, r) = \{y \in \mathbb{R}_{k,+}^n ; |x - y| < r\}$ ,  $\gamma = (\gamma_1, \dots, \gamma_k)$ ,  $\gamma_1 > 0, \dots, \gamma_k > 0$ ,  $|\gamma| = \gamma_1 + \dots + \gamma_k$ ,  $(x')^{\gamma} = x_1^{\gamma_1} \dots x_k^{\gamma_k}$ . For a measurable set  $E \subset \mathbb{R}_{k,+}^n$  suppose  $|E|_{\gamma} = \int_E (x')^{\gamma} dx$ , then  $|E(0, r)|_{\gamma} = \omega(n, k, \gamma)r^Q$ ,  $Q = n + |\gamma|$ , where

$$\omega(n, k, \gamma) = \int_{E(0,1)} (x')^{\gamma} dx = \frac{\pi^{\frac{n-k}{2}}}{2^k} \Gamma^{-1} \left( \frac{Q+2}{2} \right) \prod_{i=1}^k \Gamma \left( \frac{\gamma_i+1}{2} \right).$$

Let  $L_{p, \omega, \gamma}(\mathbb{R}_{k,+}^n)$  be the space of measurable functions on  $\mathbb{R}_{k,+}^n$  with finite norm

$$\|f\|_{L_{p, \omega, \gamma}} = \|f\|_{L_{p, \omega, \gamma}(\mathbb{R}_{k,+}^n)} = \left( \int_{\mathbb{R}_{k,+}^n} |f(x)|^p \omega^p(x) (x')^{\gamma} dx \right)^{1/p}, \quad 1 \leq p < \infty,$$

and for  $p = \infty$  the space  $L_{\infty, \omega}(\mathbb{R}_{k,+}^n)$  is defined by means of the usual modification

$$\|f\|_{L_{\infty, \omega}} = \operatorname{ess\,sup}_{x \in \mathbb{R}_{k,+}^n} \omega(x)|f(x)|.$$

Even though the  $C_{p, \gamma}$  class is well-known, for completeness, we offer the definition of  $C_{p, \gamma}$  weight functions.

**Definition 1.** *The weight function  $\omega$  belongs to the class  $C_{p, \gamma}(\mathbb{R}_{k,+}^n)$  for  $1 < p < \infty$ , if the following statement*

$$\sup_{x \in \mathbb{R}_{k,+}^n, r > 0} \left\| \frac{\omega}{|\cdot|^\gamma} \right\|_{L_{p, \gamma}(\mathbb{R}_{k,+}^n \setminus E(x, r))} \|\omega^{-1}\|_{L_{p', \gamma}(E(x, r))} < \infty$$

and

$$\sup_{x \in \mathbb{R}_{k,+}^n, r > 0} \|\omega\|_{L_{p, \gamma}(E(x, r))} \left\| \frac{1}{\omega \cdot |\cdot|^\gamma} \right\|_{L_{p', \gamma}(\mathbb{R}_{k,+}^n \setminus E(x, r))} < \infty$$

are finites.

The generalized shift operator  $T^y$  is defined by (see, for example [17], [18])

$$T^y f(x) = C_{\gamma, k} \int_0^\pi \dots \int_0^\pi f((x', y')_\beta, x'' - y'') \, d\nu(\beta),$$

where  $d\nu(\beta) = \prod_{i=1}^k \sin^{\gamma_i - 1} \beta_i \, d\beta_1 \dots d\beta_k$ ,  $x' = (x_1, \dots, x_k) \in \mathbb{R}^k$ ,

$x'' = (x_{k+1}, \dots, x_n) \in \mathbb{R}^{n-k}$ ,  $(x_i, y_i)_{\beta_i} = (x_i^2 - 2x_i y_i \cos \beta_i + y_i^2)^{\frac{1}{2}}$ ,  $1 \leq i \leq k$ ,  
 $(x', y')_\beta = ((x_1, y_1)_{\beta_1}, \dots, (x_k, y_k)_{\beta_k})$  and

$$C_{\gamma, k} = \pi^{-\frac{k}{2}} \Gamma^{-1} \left( \frac{|\gamma|}{2} \right) \prod_{i=1}^k \Gamma \left( \frac{\gamma_i + 1}{2} \right) = \frac{2^{k-1} |\gamma|}{\pi} \left( \frac{|\gamma|}{2} + 1 \right) \omega(2, k, \gamma).$$

It is well known that  $T^y$  is closely related to the Laplace-Bessel differential operator

$$\Delta_B = \sum_{i=1}^k B_i + \sum_{i=k+1}^n \frac{\partial^2}{\partial x_i^2}, \text{ where } B_i = \frac{\partial^2}{\partial x_i^2} + \frac{\gamma_i}{x_i} \frac{\partial}{\partial x_i}, \, i = 1, \dots, k.$$

**Lemma 1.** *For all  $x \in \mathbb{R}_{k,+}^n$  the following equality*

$$\int_{E(x, t)} g(y)(y')^\gamma \, dy = C_{\gamma, k}^{-1} \int_{B((x, 0), t)} g \left( \sqrt{z_1^2 + \bar{z}_1^2}, \dots, \sqrt{z_k^2 + \bar{z}_k^2}, z'' \right) \, d\nu(z, \bar{z}')$$

holds, where  $B((x, 0), t) = \{(z, \bar{z}') \in \mathbb{R}^n \times (0, \infty)^k : |(x_1 - \sqrt{z_1^2 + \bar{z}_1^2}, \dots, x_k - \sqrt{z_k^2 + \bar{z}_k^2}, x'' - z'')| < t\}$ ,  $d\nu(z, \bar{z}') = (z')^{\gamma-1} dz d\bar{z}'$ .

**Lemma 2.** For all  $x \in \mathbb{R}_{k,+}^n$ ,  $0 < \theta < \infty$ ,

$$\int_{\mathbb{R}_{k,+}^n} T^y g(x) \varphi(y) (M_\gamma \chi_{E_r})^\theta(y) (y')^\gamma dy = \int_{\mathbb{R}^n \times (0, \infty)^k} g\left(\sqrt{z_1^2 + \bar{z}_1^2}, \dots, \sqrt{z_k^2 + \bar{z}_k^2}, z''\right) \varphi(z, \bar{z}') (M_\nu \chi_{E((x,0),r)})^\theta(z, \bar{z}') d\nu(z, \bar{z}'),$$

holds where  $E((x,0),t) = \{(z, \bar{z}') \in \mathbb{R}^n \times (0, \infty)^k : |(x - z, \bar{z}')| < t\}$ .

The proofs of Lemmas 1 and 2 are done straightforwardly via the following substitutions

$$\begin{aligned} z'' &= y'', z_i = y_i \cos \alpha_i, \bar{z}_i = y_i \sin \alpha_i, \quad 0 \leq \alpha_i < \pi, \quad i = 1, \dots, k, \\ y &\in \mathbb{R}_{k,+}^n, \bar{z}' = (\bar{z}_1, \dots, \bar{z}_k), (z, \bar{z}') \in \mathbb{R}^n \times (0, \infty)^k, \quad 1 \leq k \leq n. \end{aligned}$$

**Definition 2.** [10] Let  $1 \leq p < \infty$ ,  $0 \leq \lambda \leq Q$ .  $L_{p,\lambda,\gamma}(\mathbb{R}_{k,+}^n)$ ,  $B$ -Morrey space and  $L_{p,\lambda,\omega,\gamma}(\mathbb{R}_{k,+}^n)$ , weighted  $B$ -Morrey space associated with the Laplace-Bessel differential operator defined as the space of locally integrable functions  $f(x)$ ,  $x \in \mathbb{R}_{k,+}^n$ , with the finite norm

$$\begin{aligned} \|f\|_{L_{p,\lambda,\gamma}} &= \sup_{t>0, x \in \mathbb{R}_{k,+}^n} \left( t^{-\lambda} \int_{E(0,t)} T^y |f(x)|^p (y')^\gamma dy \right)^{1/p}, \\ \|f\|_{L_{p,\lambda,\omega,\gamma}} &= \sup_{t>0, x \in \mathbb{R}_{k,+}^n} \left( t^{-\lambda} \int_{E(0,t)} T^y |f(x)|^p \omega^p(y) (y')^\gamma dy \right)^{1/p}. \end{aligned}$$

Note that

$$L_{p,0,\gamma}(\mathbb{R}_{k,+}^n) = L_{p,\gamma}(\mathbb{R}_{k,+}^n),$$

and if  $\lambda < 0$  or  $\lambda > Q$ , then  $L_{p,\lambda,\gamma}(\mathbb{R}_{k,+}^n) = \Theta$ .

### 3. Maximal Operator in $L_{p,\lambda,\omega,\gamma}(\mathbb{R}_{k,+}^n)$

Now we define the  $B$ -maximal operator by

$$M_\gamma f(x) = \sup_{r>0} |E(0,r)|_\gamma^{-1} \int_{E(0,r)} T^y |f(x)| (y')^\gamma dy.$$

Homogeneous type maximal operator defined by

$$M_\nu f(x) = \sup_{r>0} \nu(E(x,r))^{-1} \int_{E(x,r)} |f(y)| d\nu(y).$$

Also, in the work [9] it was proved:

**Proposition.** Let  $1 \leq p < \infty$  and  $\phi \in C_p(Y)$ . Then,  $M_\nu$  is bounded from  $L_{p,\phi}(Y)$  to  $L_{p,\phi}(Y)$ , where  $(Y, d, \nu)$  homogeneous type space.

The following theorem is on the boundedness of  $B$ -maximal operator  $M_\gamma$  in  $B$ -Morrey spaces.

**Theorem.** If  $f \in L_{p,\lambda,\gamma}(\mathbb{R}_{k,+}^n)$ ,  $1 < p < \infty, 0 \leq \lambda < Q$ ,  $\omega \in C_{p,\gamma}(\mathbb{R}_{k,+}^n)$ . Then,  $M_\gamma$  is bounded from  $L_{p,\lambda,\omega,\gamma}(\mathbb{R}_{k,+}^n)$  to  $L_{p,\lambda,\omega,\gamma}$ .

*Proof.* We need to introduce the maximal operator defined on a space of homogeneous type  $(Y, d, \nu)$ . By this, we mean a topological space  $Y = \mathbb{R}^n \times (0, \infty)^k$  equipped with a continuous pseudometric  $d$  and a positive measure  $\nu$  satisfying

$$\nu(E((x, \bar{x}'), 2r)) \leq C_1 \nu(E((x, \bar{x}'), r))$$

with a constant  $C_1$  independent of  $(x, \bar{x}')$  and  $r > 0$ . Here  $E((x, \bar{x}'), r) = \{(y, \bar{y}') \in Y : d((x, \bar{x}'), (y, \bar{y}')) < r\}$  ( $d\nu(y, \bar{y}')$  is given in Lemma 1),  $d((x, \bar{x}'), (y, \bar{y}')) = |(x, \bar{x}') - (y, \bar{y}')| \equiv (|x - y|^2 + |\bar{x}' - \bar{y}'|^2)^{\frac{1}{2}}$ . Let  $(Y, d, \nu)$  be a space of homogeneous type. Define

$$M_\nu \bar{f}(x, \bar{x}') = \sup_{r>0} \nu(E((x, \bar{x}'), r))^{-1} \int_{E((x, \bar{x}'), r)} |\bar{f}(y, \bar{y}')| d\nu(y),$$

where  $\bar{f}(x, \bar{x}') = f\left(\sqrt{x_1^2 + \bar{x}_1^2}, \dots, \sqrt{x_k^2 + \bar{x}_k^2}, x''\right)$ .

It can be proved that

$$M_\gamma f\left(\sqrt{z_1^2 + \bar{z}_1^2}, \dots, \sqrt{z_k^2 + \bar{z}_k^2}, z''\right) = M_\nu \bar{f}\left(\sqrt{z_1^2 + \bar{z}_1^2}, \dots, \sqrt{z_k^2 + \bar{z}_k^2}, z'', 0\right) \quad (1)$$

and

$$M_\gamma f(x) = M_\nu \bar{f}(x, 0). \quad (2)$$

Note that if  $y \in E_r = E(0, r)$ ,  $r > 0$ , then

$$M_\nu \chi_{E_r}(y) = \sup_{t>0} \frac{1}{\nu(E(y, t))} \int_{E(y, t)} \chi_{E_r}(z) d\nu(z) = \sup_{t>0} \frac{1}{\nu(E(y, t))} \int_{E(y, t) \cap E_r} d\nu(z) \leq C \sup_{t>0} \frac{1}{\nu(E(y, t))} \int_{E(y, t)} d\nu(z) \leq C.$$

The following estimate is known (see [6])

$$M_\gamma \chi_{E_r}(y) \approx \frac{r^Q}{(|y| + r)^Q}, \quad y \in \mathbb{R}_{k,+}^n, \quad r > 0.$$

It is obvious from this that  $(M_\nu \chi_{E((x,0),r)})^\theta \in C_p(Y)$  for any  $0 < \theta < 1$  is obtained.

By the Proposition and Lemma 2, taking into account  $\phi^p(y) = \omega^p(y)(M_\nu \chi_{E((x,0),r)})^\theta(y)$ ,  $\phi \in C_p(Y)$  and equalities (1) and (2), we have

$$\left( \int_{E_r} T^y (M_\gamma f(x))^p \omega^p(y) (y')^\gamma dy \right)^{\frac{1}{p}} =$$

$$\begin{aligned}
& \left( \int_Y \left( M_\nu \bar{f} \left( \sqrt{y_1^2 + \bar{y}_1^2}, \dots, \sqrt{y_k^2 + \bar{y}_k^2}, y'', 0 \right) \right)^p \times \right. \\
& \quad \left. \omega^p(y, \bar{y}') (M_\nu \chi_{E((x,0),r)})^\theta(y, \bar{y}') d\nu(y, \bar{y}') \right)^{\frac{1}{p}} = \\
& \left( \int_Y (M_\nu \bar{f}(\sqrt{y_1^2 + \bar{y}_1^2}, \dots, \sqrt{y_k^2 + \bar{y}_k^2}, y'', 0))^p \phi(y, \bar{y}') d\nu(y, \bar{y}') \right)^{\frac{1}{p}} \leq \\
& C_2 \left( \int_Y \left| \bar{f} \left( \sqrt{y_1^2 + \bar{y}_1^2}, \dots, \sqrt{y_k^2 + \bar{y}_k^2}, y'', 0 \right) \right|^p \phi(y, \bar{y}') d\nu(y, \bar{y}') \right)^{\frac{1}{p}} = \\
& \quad C_2 \left( \int_Y \left| \bar{f} \left( \sqrt{y_1^2 + \bar{y}_1^2}, \dots, \sqrt{y_k^2 + \bar{y}_k^2}, y'', 0 \right) \right|^p \times \right. \\
& \quad \left. \omega^p(y, \bar{y}') (M_\nu \chi_{E((x,0),r)})^\theta(y, \bar{y}') d\nu(y, \bar{y}') \right)^{\frac{1}{p}} = \\
& \quad C_2 \left( \int_Y \left| f \left( \sqrt{y_1^2 + \bar{y}_1^2}, \dots, \sqrt{y_k^2 + \bar{y}_k^2}, y'' \right) \right|^p \times \right. \\
& \quad \left. \omega^p(y, \bar{y}') (M_\nu \chi_{E((x,0),r)})^\theta(y, \bar{y}') d\nu(y, \bar{y}') \right)^{\frac{1}{p}} = \\
& C_2 \left( \int_{\mathbb{R}_{k,+}^n} T^y [|f|]^p(x) \omega^p(y) (M_\gamma \chi_{E_r})^\theta(y) (y')^\gamma dy \right)^{\frac{1}{p}} \leq \\
& \quad C_2 \left( \int_{E_r} T^y [|f|]^p(x) \omega^p(y) (M_\gamma \chi_{E_r})^\theta(y) (y')^\gamma dy + \right. \\
& \quad \left. \int_{\mathbb{R}_{k,+}^n \setminus E_r} T^y [|f|]^p(x) \omega^p(y) (M_\gamma \chi_{E_r})^\theta(y) (y')^\gamma dy \right)^{\frac{1}{p}} \leq \\
& \quad C_2 \left( \int_{E_r} T^y [|f|]^p(x) \omega^p(y) (y')^\gamma dy + \right. \\
& \quad \left. \sum_{j=1}^{\infty} \int_{E_{2^{j+1}r} \setminus E_{2^j r}} T^y [|f|]^p(x) \omega^p(y) (M_\gamma \chi_{E_r})^\theta(y) (y')^\gamma dy \right)^{\frac{1}{p}} \leq \\
& \quad C_2 \left( \int_{E_r} T^y [|f|]^p(x) \omega^p(y) (y')^\gamma dy + \right. \\
& \quad \left. \sum_{j=1}^{\infty} \int_{E_{2^{j+1}r} \setminus E_{2^j r}} T^y [|f|]^p(x) \omega^p(y) \frac{r^{Q\theta}}{(|y|+r)^{Q\theta}} (y')^\gamma dy \right)^{\frac{1}{p}} \leq \\
& C_3 \|f\|_{L_{p,\lambda,\omega,\gamma}} \left( r^\lambda + \sum_{j=1}^{\infty} \frac{1}{(2^j+1)^{Q\theta}} (2^{j+1}r)^\lambda \right)^{\frac{1}{p}} \leq C_4 r^{\frac{\lambda}{p}} \|f\|_{L_{p,\lambda,\omega,\gamma}}.
\end{aligned}$$

Then, we get

$$\|M_\gamma f\|_{L_{p,\lambda,\omega,\gamma}} = \sup_{x \in \mathbb{R}_{k,+}^n, t > 0} t^{-\frac{\lambda}{p}} \|T(M_\gamma f(x))\|_{L_{p,\omega,\gamma}(E_t)} \leq C_4 \|f\|_{L_{p,\lambda,\omega,\gamma}}.$$



**Corollary.** *If  $1 < p < \infty$ ,  $f \in L_{p,\gamma}(\mathbb{R}_{k,+}^n)$ ,  $\omega \in C_{p,\gamma}(\mathbb{R}_{k,+}^n)$ . Then,  $M_\gamma$  is bounded from  $L_{p,\omega,\gamma}(\mathbb{R}_{k,+}^n)$  to  $L_{p,\omega,\gamma}$ .*

## References

1. Abasova G.A., Aliyeva L.R., Hasanov J.J., Shirinova E.S. Necessary and sufficient conditions for the boundedness of comutators of  $B$ -Riesz potentials in Lebegues spaces. *J. Contemp. Appl. Math.*, 2016, **6** (2), pp. 18-32.
2. Aliev I.A., Bayrakci S. On inversion of  $B$ -elliptic potentials by the method of Balakrishnan-Rubin. *Fract. Calc. Appl. Anal.*, 1998, **1** (4), pp. 365-384.
3. Aliyeva L.R., Esen Almali S., Safarov Z.V. Maximal operator in weighted Morrey spaces, associated with the Laplace-Bessel differential operator. *Baku Math. J.*, 2023, **2** (2), pp. 176-183.
4. Ayazoglu R., Hasanov J.J. On the boundedness of a  $B$ -Riesz potential in the generalized weighted  $B$ -Morrey spaces. *Georgian Math. J.*, 2016, **23** (2), pp. 143-155.
5. Aykol C., Hasanov J.J. On the boundedness of  $B$ -maximal commutators, commutators of  $B$ -Riesz potentials and  $B$ -singular integral operators in modified  $B$ -Morrey spaces. *Acta Sci. Math.*, 2020, **86** (3-4), pp. 521-547.
6. Burenkov V.I., Guliyev H.V. Necessary and sufficient conditions for boundedness of the maximal operator in the local Morrey-type spaces. *Studia Math.*, 2004, **163** (2), pp. 157-176.
7. Chiarenza F., Frasca M. Morrey spaces and Hardy-Littlewood maximal function. *Rend. Math.*, 1987, **7**, pp. 273-279.
8. Gadjiev A.D., Aliev I.A. On classes of operators of potential types, generated by a generalized shift. *Reports of Enlarged Session of the Seminars of I.N. Vekua Inst. Appl. Math.*, 1988, **3** (2), pp. 21-24 (in Russian).
9. García-Cuerva J., Rubio de Francia J.L. *Weighted Norm Inequalities and Related Topics*, **116**. North-Holland Publ. Co., Amsterdam, 1985.
10. Guliev V.S. Sobolev's theorem for the anisotropic Riesz-Bessel potential in Morrey-Bessel spaces. *Dokl. Akad. Nauk*, 1999, **367** (2), pp. 155-156 (in Russian).
11. Guliev V.S. On maximal function and fractional integral, associated with the Bessel differential operator. *Math. Inequal. Appl.*, 2003, **6** (2), pp. 317-330.
12. Guliyev V.S., Hasanov J.J. Sobolev-Morrey type inequality for Riesz potentials, associated with the Laplace-Bessel differential operator. *Fract. Calc. Appl. Anal.*, 2006, **9** (1), pp. 17-32.
13. Guliyev V.S., Hasanov J.J. Necessary and sufficient conditions for the boundedness of  $B$ -Riesz potential in the  $B$ -Morrey spaces. *J. Math. Anal. Appl.*, 2008, **347**, pp. 113-122.

14. Hasanov J.J. A note on anisotropic potentials, associated with the Laplace-Bessel differential operator. *Oper. Matrices*, 2008, **2** (4), pp. 465-481.
15. Hasanov J.J., Safarov Z.V. Hardy-Littlewood-Stein-Weiss theorems for Riesz potentials in modified Morrey spaces. *Baku Math. J.*, 2022, **1** (1), pp. 37-46.
16. Hasanov J.J., Zeren Y. On limiting case of the Sobolev theorem for  $B$ -Riesz potential in  $B$ -Morrey spaces. *Arab J. Math. Sci.*, 2007, **13**, pp. 27-38.
17. Kipriyanov I.A. Fourier-Bessel transformations and embedding theorems. *Trudy Math. Inst. Steklov*, 1967, **89**, pp. 130-213 (in Russian).
18. Levitan B.M. Bessel function expansions in series and Fourier integrals. *Uspekhi Mat. Nauk*, 1951, **6** (2(42)), pp. 102-143 (in Russian).
19. Lyakhov L.N. Multipliers of the mixed Fourier-Bessel transform. *Proc. Steklov Inst. Math.*, 1996, **214**, pp. 227-242.
20. Morrey C.B. On the solutions of quasi-linear elliptic partial differential equations. *Trans. Amer. Math. Soc.*, 1938, **43**, pp. 126-166.
21. Muckenhoupt B., Stein E.M. Classical expansions and their relation to conjugate harmonic functions. *Trans. Amer. Math. Soc.*, 1965, **118**, pp. 17-92.
22. Şerbetçi A., Ekincioglu I. Boundedness of Riesz potential generated by generalized shift operator on  $Ba$  spaces. *Czechoslovak Math. J.*, 2004, **54** (3), pp. 579-589.
23. Shishkina E.L. Hyperbolic Riesz  $B$ -potential and solution of an iterated non-homogeneous  $B$ -hyperbolic equation. *Lobachevskii J. Math.*, 2020, **41** (5), pp. 895-916.
24. Stempak K. Almost everywhere summability of Laguerre series. *Studia Math.*, 1991, **100** (2), pp. 129-147.
25. Triméche K. Inversion of the Lions transmutation operators using generalized wavelets. *Appl. Comput. Harmon. Anal.*, 1997, **4** (1), pp. 97-112.