

FORMULAS FOR THE SUMS OF SOME CONDITIONALLY CONVERGENT SERIES

B.D. BARMAK

Received: 12.01.2026 / Revised: 09.02.2026 / Accepted: 16.02.2026

Abstract. *In this paper, we find formulas for the sums of two families of conditionally convergent series containing $p(\geq 1)$ arbitrary real parameters, in terms of elementary functions. These families, when the parameters are chosen appropriately, coincide with the formulas for $\ln 2$ and $\pi/4$ when $p = 1$, and with the formulas known to Newton when $p = 2$ and to Euler when $p = 2, 4$.*

Keywords: conditionally convergent series, Newton-Euler summation formulas, elementary functions

Mathematics Subject Classification (2020): 26A09, 40C15

1. Introduction

Consider two divergent series

$$1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots \quad \text{and} \quad 1 + \frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \dots$$

Let p be a natural number. Without rearranging the terms of these series, we change their signs so that p positive terms are followed by the same number of negative terms, then p positive terms again, and so on. This results in conditionally convergent series

$$S_{1,p} := 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{p} - \frac{1}{p+1} - \frac{1}{p+2} - \dots - \frac{1}{2p} + \frac{1}{2p+1} + \dots + \frac{1}{3p} - \dots,$$

$$S_{2,p} := 1 + \frac{1}{3} + \frac{1}{5} + \dots + \frac{1}{2p-1} - \frac{1}{2p+1} - \frac{1}{2p+3} - \dots - \frac{1}{4p-1} + \frac{1}{4p+1} + \dots + \frac{1}{6p-1} - \dots$$

These series are special cases of a conditionally convergent series

$$S_{b,p}(a_0, a_1, \dots, a_{p-1}) := \sum_{n=0}^{+\infty} (-1)^n \left(\frac{a_0}{bpn+1} + \dots + \frac{a_{p-1}}{bpn+b(p-1)+1} \right), \quad (1)$$

Bella D. Barmak

Lomonosov Moscow State University, Moscow, Russia;

Northern (Arctic) Federal University named after M.V. Lomonosov, Arkhangelsk, Russia

E-mail: barmakbella@mail.ru

where a_0, a_1, \dots, a_{p-1} are given real numbers, and the parameter b is equal to 1 or 2 when $a_0 = \dots = a_{p-1} = 1$.

The classical formulas $S_{1,1} = \ln 2$ and $S_{2,1} = \pi/4$ are well known. They are obtained from the Taylor series expansions of the functions $\ln(1+x)$ and $\operatorname{arctg} x$, respectively, and the limit transition in the resulting series as $x \rightarrow 1$. The very interesting history of the discovery of these formulas is described in [5].

In this paper, we present a method (see Section 2) that allows us to find formulas for the sum $S_{b,p}(a_0, a_1, \dots, a_{p-1})$ in terms of elementary functions. In particular, in Section 3, we prove the following theorem.

Theorem. *Let $p \in \mathbb{N}$ and $p \geq 2$. Then the following equalities hold*

$$S_{1,p} = \frac{\ln 2}{p} + \frac{\pi}{p} \sum_{j=0}^{[(p-2)/2]} \left(1 - \frac{1+2j}{p}\right) \operatorname{ctg} \left(\frac{\pi + 2\pi j}{2p}\right) \quad (2)$$

and

$$S_{2,p} = \frac{\pi}{2p} \left(\frac{\varepsilon_p}{2} + \sum_{j=0}^{[(p-2)/2]} \frac{1}{\sin \frac{\pi + 2\pi j}{2p}} \right), \quad (3)$$

where $[x]$ is the integer part of x , and $\varepsilon_p = 0$ if p is an even number, and $\varepsilon_p = 1$ if p is an odd number.

Letting $p = 2, 3, 4, 5$ and 6 in the formulas (2) and (3), we obtain the following identities

$$S_{1,2} = \frac{\ln 2}{2} + \frac{\pi}{4}, \quad S_{1,3} = \frac{\ln 2}{3} + \frac{2\pi}{3\sqrt{3}}, \quad S_{1,4} = \frac{\ln 2}{4} + \frac{(2\sqrt{2} + 1)\pi}{8},$$

$$S_{1,5} = \frac{\ln 2}{5} + \frac{2\sqrt{25 + 10\sqrt{5}} \cdot \pi}{25}, \quad S_{1,6} = \frac{\ln 2}{6} + \frac{(15 + 4\sqrt{3})\pi}{36},$$

$$S_{2,2} = \frac{\pi}{2\sqrt{2}}, \quad S_{2,3} = \frac{5\pi}{12}, \quad S_{2,4} = \frac{\sqrt{2 + \sqrt{2}} \cdot \pi}{4},$$

$$S_{2,5} = \frac{(4\sqrt{5} + 1)\pi}{20}, \quad S_{2,6} = \frac{\sqrt{2}(1 + 2\sqrt{3})\pi}{12}$$

accordingly.

In conclusion of this paragraph, we note that some of these formulas are well known, for example, the formulas for $S_{1,2}$ and $S_{2,3}$ are given in [4, clause 5.1.2, formula 5, and clause 5.1.4, formula 5], the formula for $S_{2,2}$ belongs to Newton (see, for example, [1, Chapter 1, problem No 52(b)] and [3]), and the formula for $S_{2,4}$ belongs to Euler (see, for example, [2, p. 180]).

2. The Method of Calculating the Amount

$S_{b,p}(a_0, a_1, \dots, a_{p-1})$

2. 1.

Let the numbers $b, p, a_0, a_1, \dots, a_{p-1}$ be the same as in Section 1. This section provides a method for finding formulas for the sums of conditionally convergent series (1).

Consider the power series

$$f_{b,p}(x) (\equiv f_{b,p}(a_0, a_1, \dots, a_{p-1}; x)) := \sum_{n=0}^{+\infty} (-1)^n \left(\frac{a_0 x^{bpn+1}}{bpn+1} + \dots + \frac{a_{p-1} x^{bpn+b(p-1)+1}}{bpn+b(p-1)+1} \right).$$

It is easy to see that this series converges for $|x| < 1$ and allows for pointwise differentiation. Therefore

$$f'_{b,p}(x) = \sum_{n=0}^{+\infty} (-x^{bp})^n (a_0 + \dots + a_{p-1} x^{b(p-1)}) = \frac{P_{b,p}(x)}{1+x^{bp}},$$

where $P_{b,p}(x) := a_0 + a_1 x^b + \dots + a_{p-1} x^{b(p-1)}$. Given that $f_{b,p}(0) = 0$ and integrating (2) over the interval $[0; x]$, we find that

$$f_{b,p}(x) = \int_0^x \frac{P_{b,p}(t)}{1+t^{bp}} dt. \quad (4)$$

Next, let $x_0, x_1, \dots, x_{bp-1}$ denote the roots of the equation

$$1+x^{bp} = 0 \quad (5)$$

and by expanding the integrand into simple fractions, we find that

$$\frac{P_{b,p}(t)}{1+t^{bp}} = \frac{A_0}{t-x_0} + \frac{A_1}{t-x_1} + \dots + \frac{A_{bp-1}}{t-x_{bp-1}}, \quad (6)$$

where the unknown coefficients A_0, \dots, A_{bp-1} are given by the formula

$$A_j = \lim_{t \rightarrow x_j} \frac{P_{b,p}(t)(t-x_j)}{1+t^{bp}} = -\frac{x_j}{bp} P_{b,p}(x_j), \quad j = 0, 1, \dots, bp-1. \quad (7)$$

2. 2.

First, consider the case $b = 1$ and $p = 2k$, $k = 1, 2, \dots$. In this case, the roots of the equation (5) are found by the formulas $x_j = e^{i\pi(1+2j)/(2k)}$, $j = 0, 1, \dots, 2k-1$. From these equalities and the realness of the numbers a_0, a_1, \dots, a_{p-1} , it follows that for $j = 0, \dots, k-1$, the equalities $x_{2k-1-j} = \bar{x}_j$ and $A_{2k-1-j} = \bar{A}_j$. Therefore, by

grouping the corresponding terms in (6), taking into account the obtained result in (4) and integrating, we arrive at the validity of the relation

$$f_{1,2k}(x) = \sum_{j=0}^{k-1} \operatorname{Re} A_j \cdot \ln(x^2 - 2\operatorname{Re} x_j \cdot x + 1) - 2 \sum_{j=0}^{k-1} \operatorname{Im} A_j \cdot \left(\operatorname{arctg} \left(\frac{x - \operatorname{Re} x_j}{\operatorname{Im} x_j} \right) + \frac{\pi}{2} - \frac{\pi + 2\pi j}{2k} \right). \quad (8)$$

Taking the limit of both sides of this equality as $x \rightarrow 1$ and using Abel's second theorem, we find that

$$S_{1,2k}(a_0, a_1, \dots, a_{2k-1}) = f_{1,2k}(a_0, a_1, \dots, a_{2k-1}; 1). \quad (9)$$

Now consider the case when $b = 1$ and $p = 2k + 1$, $k = 1, 2, \dots$. In this case, the roots of equation (5) are given by the formula

$$x_j = e^{\frac{i\pi(1+2j)}{2k+1}}, \quad j = 0, 1, \dots, 2k. \quad (10)$$

Using arguments similar to those above, we can prove that the following equalities are true

$$f_{1,2k+1}(x) = \frac{P_{1,2k+1}(-1)}{2k+1} \ln(x+1) + \sum_{j=0}^{k-1} \operatorname{Re} A_j \ln(x^2 - 2\operatorname{Re} x_j \cdot x + 1) - 2 \sum_{j=0}^{k-1} \operatorname{Im} A_j \left(\operatorname{arctg} \left(\frac{x - \operatorname{Re} x_j}{\operatorname{Im} x_j} \right) + \frac{\pi}{2} - \frac{\pi + 2\pi j}{2k+1} \right), \quad (11)$$

$$S_{1,2k+1}(a_0, a_1, \dots, a_{2k}) = f_{1,2k+1}(a_0, a_1, \dots, a_{2k}; 1), \quad (12)$$

where A_j is calculated using formula (7), and x_j is calculated using formula (10).

2. 3.

If $b = 2$, then repeating the reasoning above, we conclude that

$$f_{2,p}(x) = \sum_{j=0}^{p-1} \operatorname{Re} A_j \ln(x^2 - 2\operatorname{Re} x_j \cdot x + 1) - 2 \sum_{j=0}^{p-1} \operatorname{Im} A_j \left(\operatorname{arctg} \left(\frac{x - \operatorname{Re} x_j}{\operatorname{Im} x_j} \right) + \frac{\pi}{2} - \frac{\pi + 2\pi j}{2p} \right), \quad (13)$$

where x_j are defined as the roots of equation (5), and A_j are calculated using formula (7) with $b = 2$, and

$$S_{2,p}(a_0, a_1, \dots, a_{p-1}) = f_{2,p}(a_0, a_1, \dots, a_{p-1}; 1). \quad (14)$$

Formulas (8), (11), (13) allow us to find the sums of power series $f_{b,p}(x)$, and formulas (9), (12), (14) - sums of conditionally convergent numerical series $S_{b,p}(a_0, a_1, \dots, a_{p-1})$ in terms of elementary functions.

3. Proof of Theorem

3. 1.

Let's move on to the proof of the formula (2). When $b = 1$ and $a_0 = a_1 = \dots = a_{p-1} = 1$, it follows from (7) that

$$A_j = \frac{2x_j}{p(x_j - 1)}, \quad j = 0, 1, \dots, p - 1.$$

From this equality, it can be deduced that

$$\operatorname{Re} A_j = \frac{1}{p}, \quad \operatorname{Im} A_j = -\frac{1}{p} \operatorname{ctg} \left(\frac{\pi + 2\pi j}{2p} \right), \quad j = 0, \dots, p - 1.$$

Thus, in the case $p = 2k$, $k = 1, 2, \dots$, the equality (9) can be written as

$$S_{1, 2k} = \frac{\ln 2}{2} + \frac{1}{2k} \sum_{j=0}^{k-1} \ln \left(1 - \cos \frac{\pi + 2\pi j}{2k} \right) + \frac{1}{2k} \sum_{j=0}^{k-1} \operatorname{ctg} \left(\frac{\pi + 2\pi j}{4k} \right) \cdot \left(\pi - \frac{\pi + 2\pi j}{2k} \right).$$

Using the formula

$$\prod_{j=0}^{\lfloor \frac{n-1}{2} \rfloor} \sin \frac{\pi j}{n} = \sqrt{2^{1-n} n}$$

(see, for example, [1, p. 6.1.2, formula 3]), we transform the first sum

$$\sum_{j=0}^{k-1} \ln \left(1 - \cos \frac{\pi + 2\pi j}{2k} \right) = k \ln 2 + 2 \left(\ln \left(\prod_{j=0}^{\lfloor \frac{4k-1}{2} \rfloor} \sin \frac{\pi j}{4k} \right) - \ln \left(\prod_{j=0}^{\lfloor \frac{2k-1}{2} \rfloor} \sin \frac{\pi j}{2k} \right) \right) = (1-k) \ln 2.$$

Thus, we finally get that

$$S_{1, 2k} = \frac{\ln 2}{2k} + \frac{\pi}{2k} \sum_{j=0}^{k-1} \left(1 - \frac{1 + 2j}{2k} \right) \operatorname{ctg} \left(\frac{\pi + 2\pi j}{4k} \right).$$

If $p = 2k + 1$, $k = 1, 2, \dots$, then

$$S_{1, 2k+1} = \frac{k+1}{2k+1} \ln 2 + \frac{1}{2k+1} \sum_{j=0}^{k-1} \left[\ln \left(1 - \cos \frac{\pi + 2\pi j}{2k+1} \right) + \left(\pi - \frac{\pi + 2\pi j}{2k+1} \right) \operatorname{ctg} \frac{\pi + 2\pi j}{2(2k+1)} \right],$$

where, as calculations show,

$$\sum_{j=0}^{k-1} \ln \left(1 - \cos \frac{\pi + 2\pi j}{2k+1} \right) = -k \ln 2.$$

Therefore, formula (2) is also valid for $p = 2k + 1$.

3. 2.

In the case $b = 2$ and $a_0 = a_1 = \dots = a_{p-1} = 1$, it follows from formula (7) that

$$A_j = \frac{x_j}{p(x_j^2 - 1)}, \quad j = 0, 1, \dots, 2p - 1.$$

From this equality, it can be deduced that

$$\operatorname{Re} A_j = 0, \quad \operatorname{Im} A_j = -\frac{1}{2p} \cdot \frac{1}{\sin \frac{\pi + 2\pi j}{2p}}.$$

Considering these equalities in (14), we arrive at the identity

$$S_{2,p} = \frac{\pi}{4p^2} \sum_{j=0}^{p-1} \frac{2(p-j) - 1}{\sin \frac{\pi + 2\pi j}{2p}},$$

from which, in turn, the equality (3) follows.

Acknowledgements The author expresses deep gratitude to his supervisor, Professor K. A. Mirzoev, for setting the task and for his constant attention to the work.

The work was carried out with financial support from the Russian Science Foundation under research project No. 24-21-00128.

References

1. Askey R., Roy R., Andrews G.E. *Special Functions*. Publ. House Moscow Center Continuous Math. Edu., Moscow, 2013 (in Russian).
2. Euler L. *Introduction to Infinitesimal Analysis*, **1**, 2nd edition. State Publ. House Phys. Math., Moscow, 1961 (in Russian).
3. Newton I. *The Correspondence of Isaac Newton*, **2**. Cambridge Univ. Press, Cambridge, 1960, pp. 1676–1687.
4. Prudnikov A.P., Brychkov Yu.A., Marichev O.I. *Integrals and Series*, **1**. *Elementary Functions*, 2nd edition, revised. Fizmatlit, Moscow, 2002 (in Russian).
5. Roy R. The discovery of the series formula for π by Leibniz, Gregory and Nilakantha. *Math. Mag.*, 1990, **63** (5), pp. 291–306.